



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 7/04, C07K 14/135	A1	(11) International Publication Number: WO 97/12032 (43) International Publication Date: 3 April 1997 (03.04.97)
(21) International Application Number: PCT/US96/15524 (22) International Filing Date: 27 September 1996 (27.09.96) (30) Priority Data: 60/007,083 27 September 1995 (27.09.95) US (71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as represented by THE DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; Box OTT, Bethesda, MD 20892 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): COLLINS, Peter, L. [US/US]; #401 12304 Village Square Terrace, Rockville, MD 20852 (US). (74) Agents: PARMELEE, Steven, W. et al.; Townsend and Townsend and Crew L.L.P., 8th floor, Two Embarcadero Center, San Francisco, CA 94111 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PRODUCTION OF INFECTIOUS RESPIRATORY SYNCYTIAL VIRUS FROM CLONED NUCLEOTIDE SEQUENCES		
(57) Abstract <p>Isolated polynucleotide molecules provide RSV genome and antigenomes, including that of human, bovine or murine RSV or RSV-like viruses, and chimera thereof. The recombinant genome or antigenome can be expressed with a nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large (L) polymerase protein, and an RNA polymerase elongation factor to produce isolated infectious RSV particles. The recombinant RSV genome and antigenome can be modified to produce desired phenotypic changes, such as attenuated viruses for vaccine use.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic			SE	Sweden
CG	Congo	KR	Republic of Korea	SG	Singapore
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LR	Liberia	SZ	Swaziland
CS	Czechoslovakia	LT	Lithuania	TD	Chad
CZ	Czech Republic	LU	Luxembourg	TG	Togo
DE	Germany	LV	Latvia	TJ	Tajikistan
DK	Denmark	MC	Monaco	TT	Trinidad and Tobago
EE	Estonia	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	UG	Uganda
FI	Finland	ML	Mali	US	United States of America
FR	France	MN	Mongolia	UZ	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Viet Nam

PRODUCTION OF INFECTIOUS RESPIRATORY SYNCYTIAL
VIRUS FROM CLONED NUCLEOTIDE SEQUENCES

Background of the Invention

Human respiratory syncytial virus (RSV) is the most important pediatric respiratory pathogen worldwide. This ubiquitous, highly infectious agent emerges each year in seasonal epidemics. Nearly everyone is infected at least once within the first two years of life. RSV disease is responsible for considerable morbidity and mortality, especially in the very young; in the United States it causes an estimated 91,000 hospitalizations and 4500 deaths annually, and its impact is much greater in less affluent countries. RSV also has come to be recognized as an important agent of disease of immunocompromised adults and of the elderly.

Resistance to RSV reinfection induced by natural infection is incomplete but increases incrementally with repeated exposure. Thus, RSV can infect multiple times during childhood and life, but serious disease usually is limited to the first and sometimes second infections of life. The minimum goal of RSV immunoprophylaxis is to induce sufficient resistance to prevent serious disease associated with the initial infections.

A number of attenuated RSV strains were developed and evaluated as vaccines during the 1960's and 70's, but they were found to be either over- or under-attenuated, and in some cases exhibited genetic instability, as is common for single-stranded RNA viruses. Current strategies under investigation for RSV vaccine development are principally the parenteral administration of purified viral antigen or the development of live attenuated RSV for intranasal administration. The

intranasal route provides direct stimulation of local immunity. It also partially abrogates the immunosuppressive effects of RSV-specific maternally derived serum antibodies, which typically are found in the very young. The parenteral administration of inactivated RSV or purified RSV antigen in experimental animals appears to be associated with enhanced immunopathology upon subsequent virus challenge, similar to the enhanced RSV disease associated with a formalin-inactivated vaccine evaluated in the 1960's. But this effect has never been observed with RSV infection of the respiratory tract, suggesting that live attenuated viruses have an important advantage in safety. To date, however, there is no approved vaccine or highly effective antiviral therapy for RSV.

Research efforts to produce a suitable RSV vaccine are impeded by poor viral growth in tissue culture, a lengthy replication cycle, virion instability, a negative-sense RNA genome, and a complex genome organization and gene products. RSV is a member of the pneumovirus genus of the paramyxovirus family, and its genome of single-stranded negative-sense RNA of 15,222 nucleotides has been sequenced completely for wild-type strain A2 virus as well as for an attenuated derivative thereof.

Some aspects of RNA synthesis by RSV appear to follow the general pattern of nonsegmented negative strand viruses. The genome template is tightly encapsidated with the major nucleocapsid (N) protein and is associated with the phosphoprotein (P) and large (L) polymerase subunit protein. Transcription begins at the 3' extragenic leader region and proceeds along the entire length by a sequential, stop-start mechanism guided by short template signals flanking the genes. This yields at least ten major species of mRNA which encode at least ten major proteins. RNA replication occurs by a switch to the synthesis of a full length positive-sense "antigenome" which also is tightly encapsidated and serves as the template for the synthesis of progeny genome.

The viral genomic RNA of negative-strand viruses is not infectious alone as free RNA. In virions or intracellularly, viral RNA is always found tightly encapsidated in a ribonucleoprotein core. This nucleocapsid contains the viral proteins necessary for transcription and replication and has long been regarded as the minimum unit of infectivity (Brown et al., J. Virol. 1: 368-373 (1967)). Thus, it has been recognized that the generation of biologically active synthetic viral RNA from cDNA will require complementation by viral protein, leading to the assembly of functional nucleocapsids (Collins et al., Proc. Natl. Acad. Sci. USA 88: 9663-9667 (1991), and Collins et al., Virology 195: 252-256 (1993)). The ability to produce live RSV from cDNA is of particular importance because it would permit the introduction of specific engineered changes, including attenuating mutations, into the genome of infectious virus in an effort to produce safe and effective RSV vaccines.

Short, internally-deleted analogs of genome or antigenome RNA ("minigenomes") have been shown to participate in transcription and replication when synthesized intracellularly in the presence of the appropriate viral proteins. For two rhabdoviruses, rabies and vesicular stomatitis viruses, infectious virus has been produced by coexpression of a complete cDNA-encoded antigenome RNA in the presence of the N, P and L proteins (Schnell et al., EMBO J. 13: 4195-4203 (1994) and Lawson et al., Proc. Natl. Acad. Sci. USA 92: 4477-4481 (1995)).

RSV possesses a number of properties which distinguishes it and other members of the genus *Pneumovirus* from the better characterized paramyxoviruses of the genera *Paramyxovirus*, *Rubulavirus* and *Morbillivirus*. These differences include a greater number of mRNAs, an unusual gene order at the 3' end of the genome, species-to-species variability in the order of the glycoprotein and M2 genes, a greater diversity in intergenic regions, an attachment protein that exhibits mucin-like characteristics, extensive strain-to-strain sequence diversity, and several proteins not found in

any or most of the other nonsegmented negative strand RNA viruses.

RSV remains the most common cause of severe viral lower respiratory tract disease in infants and children.

5 Consequently, an urgent need remains for the ability to engineer a safe and effective vaccine that is able to prevent the serious illness in this population that often requires hospitalization. Quite surprisingly, the present invention fulfills this and other related needs by providing methods for
10 introducing defined, predetermined changes into infectious RSV.

Summary of the invention

15

The present invention provides an isolated infectious RSV particle which comprises a recombinant RSV genome or antigenome, a nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large (L) polymerase
20 protein, and an RNA polymerase elongation factor. The RNA polymerase elongation factor can be M2(ORF1) of RSV. The isolated infectious RSV particle can be a viral or subviral particle. The isolated infectious RSV virus may be a human RSV, a bovine or murine RSV, or the genome or antigenome can
25 be a chimera of two or more different RSV genomes, such as having nucleotide segments from human and bovine RSV.

In other embodiments the invention provides a method for producing an infectious RSV particle from one or more isolated polynucleotide molecules encoding an RSV. An
30 expression vector which comprises an isolated polynucleotide molecule encoding a RSV genome or antigenome and an expression vector which comprises one or more isolated polynucleotide molecules that encodes N, P, L and RNA polymerase elongation factor proteins are coexpressed in a cell or cell-free lysate,
35 thereby producing an infectious RSV particle. The RSV genome or antigenome and the N, P, L and RNA polymerase elongation factor proteins can be coexpressed by the same or different

expression vectors. In some instances the N, P, L and RNA polymerase elongation factor proteins are each encoded on different expression vectors. The polynucleotide molecule encoding the RSV genome or antigenome is from a human, bovine or murine RSV sequence, and can be a chimera of a human RSV strain sequence and at least one non-human RSV sequence, or can encode the genome or antigenome of a wild-type RSV strain. The RSV genome or antigenome can be modified from a wild-type RSV strain by a nucleotide insertion, rearrangement, deletion or substitution, so as to encode a phenotypic alteration such as one that results in attenuation, temperature-sensitivity, cold-adaptation, small plaque size, host range restriction, or a change in an immunogenic epitope of RSV. The polynucleotide can encode a genome or antigenome of a nonhuman RSV virus, or can be a chimera of a nonhuman RSV and at least one other RSV or human or nonhuman origin. The polynucleotide molecule encoding the genome or antigenome can also be modified to include a nucleotide sequence that encodes a cytokine, a T-helper epitope, a G protein of a different RSV subgroup, a restriction site marker, or a protein of a microbial pathogen (e.g., virus, bacterium or fungus) capable of eliciting a protective immune response in the intended host.

In other embodiments the invention provides a cell or cell-free lysate containing an expression vector which comprises an isolated polynucleotide molecule encoding a RSV genome or antigenome and an expression vector which comprises one or more isolated polynucleotide molecules that encodes N, P, L and RNA polymerase elongation factor proteins of RSV. Upon expression the genome or antigenome and N, P, L, and RNA polymerase elongation factor proteins combine to produce an infectious RSV particle, such as viral or subviral particle.

In another aspect the invention provides an isolated polynucleotide molecule which comprises an operably linked transcriptional promoter, a polynucleotide sequence encoding an RSV genome or antigenome, and a transcriptional terminator. The RSV genome or antigenome can be a human RSV

sequence and modified versions thereof, such as that exemplified in SEQ ID NO:1 (which depicts the 5' to 3' positive-sense sequence whereas the genome itself is negative-sense). The polynucleotide can also encode a genome or antigenome of a nonhuman RSV virus, or encode a chimera of a nonhuman RSV and at least one other RSV of human or nonhuman origin.

10

Brief Description of the Drawings

Fig. 1A and 1B show the construction of cDNA encoding RSV antigenome RNA, where Fig. 1A shows the structures of the cDNA and the encoded antigenome RNA (not to scale). The diagram of the antigenome includes the following features: the 5'-terminal nonviral G triplet contributed by the T7 promoter, the four sequence markers at positions 1099 (which adds one nt to the length), 1139, 5611, and 7559 (numbering referring to the first base of the new restriction site), the ribozyme and tandem T7 terminators, and the single nonviral 3'-phosphorylated U residue contributed to the 3' end by ribozyme cleavage (the site of cleavage is indicated with an arrow). Cloned cDNA segments representing in aggregate the complete antigenome are also shown. The box illustrates the removal of the BamHI site, a modification that facilitated assembly: the naturally occurring BamHI-SalI fragment (the BamHI site is shown in the top line in positive sense, underlined) was replaced with a PCR-generated BglII-SalI fragment (the BglII site is shown in the bottom line, underlined; its 4-nt sticky end [shown in italics] is compatible with that of BamHI). This resulted in a single nt change (middle line, underlined) which was silent at the amino acid level.

Fig. 1B shows the sequence markers contained in the cDNA-encoded antigenome RNA, where sequences are positive sense and numbered relative to the first nt of the leader region complement as 1; identities between strains A2 and

18537, representing subgroups A and B, respectively, are indicated with dots; sequences representing restriction sites in the cDNA are underlined; gene-start (GS) and gene-end (GE) transcription signals are boxed; the initiation codon of the N translational open reading frame at position 1141 is italicized, and the sequence markers are shown underneath each sequence. In the top sequence, a single C residue was inserted at position 1099 to create an AflIII site in the NSII-N intergenic region, and the AG at positions 1139 and 1140 immediately upstream of the N translational open reading frame were replaced with CC to create a new NcoI site. In the middle sequence, substitution of G and U at positions 5612 and 5616, respectively, created a new StuI site in the G-F intergenic region. In the bottom sequence, a C replacement at position 7560 created a new SphI site in the F-M2 intergenic region.

Fig. 2 shows construction of D46/1024CAT cDNA encoding an RSV antigenome containing the CAT ORF flanked by RSV transcription signals (not to scale, RSV-specific segments are shown as filled boxes and CAT sequence as an open box). The source of the CAT gene transcription cassette was RSV-CAT minigenome cDNA 6196 (diagram at top). The RSV-CAT minigenome contains the leader region, gene-start (GS) and gene-end (GE) signals, noncoding (NC) RSV gene sequences, and the CAT ORF, with *XmaI* restriction endonuclease sites preceding the GS signal and following the GE signal. The nucleotide lengths of these elements are indicated, and the sequences (positive-sense) surrounding the *XmaI* sites are shown above the diagram. A 8-nucleotide *XmaI* linker was inserted into *StuI* site of the parental plasmid D46 to construct the plasmid D46/1024. The *XmaI* - *XmaI* fragment of the plasmid 6196 was inserted into the plasmid D46/1024 to construct the plasmid D46/1024CAT. The RNA encoded by the D46 cDNA is shown at the bottom, including the three 5'-terminal nonviral G residues contributed by the T7 promoter and the 3'-terminal phosphorylated U residue contributed by cleavage of the hammerhead ribozyme; the

nucleotide lengths do not include these nonviral nucleotides. The L gene is drawn offset to indicate the gene overlap.

5 Description of the Specific Embodiments

The present invention provides the production of infectious RSV from cDNA. Infectious RSV is produced by the intracellular coexpression of a cDNA that encodes the RSV genome or antigenome RNA, together with those viral proteins necessary to generate a transcribing, replicating nucleocapsid, preferably one or more sequences that encode major nucleocapsid (N or NP) protein, nucleocapsid phosphoprotein (P), large (L) polymerase protein, and an M2(ORF1) protein. Infectious RSV particles are produced by the recombinant system. The recombinant production system permits the introduction of defined changes into infectious RSV, which changes are useful in a wide variety of applications such as: the development of live attenuated vaccine strains bearing predetermined, defined attenuating mutations; analysis of RSV molecular biology and pathogenesis using, e.g., defined mutations to alter functions or expression of RSV proteins; improvement in the growth in culture; identification of attenuating mutations in existing or future vaccine strains by distinguishing between silent incidental mutations versus those responsible for phenotype differences; production of modified vaccine virus to accommodate antigenic drift; enhancement of vaccine immunogenicity; ablation of epitopes associated with undesirable immunopathology; insertion of foreign genes, in whole or in part, encoding protective antigens to generate RSV strains capable of inducing immunity to both RSV and the virus or agent from which the protective antigen was derived; insertion of foreign genes, in whole or in part, encoding modulators of the immune system such as cytokines or T cell epitopes, to enhance the immunogenicity of the vaccine virus; etc.

According to the present invention cDNA encoding a RSV genome or antigenome is constructed for intracellular or in vitro coexpression with the necessary viral proteins to form infectious RSV. By "RSV antigenome" is meant an isolated positive-sense polynucleotide molecule which serves as the template for the synthesis of progeny RSV genome. Preferably a cDNA is constructed which is a positive-sense version of the RSV genome, corresponding to the replicative intermediate RNA, or antigenome, so as to minimize the possibility of hybridizing with positive-sense transcripts of the complementing sequences that encode proteins necessary to generate a transcribing, replicating nucleocapsid, i.e., sequences that encode N, P, L and M2(ORF1) protein. In an RSV minigenome system, genome and antigenome were equally active in rescue, whether complemented by RSV or by plasmids, indicating that either genome or antigenome can be used and thus the choice can be made on methodologic or other grounds.

A native RSV genome typically comprises a negative-sense polynucleotide molecule which, through complementary viral mRNAs, encodes eleven species of viral proteins, i.e., the nonstructural species NS1 and NS2, N, P, matrix (M), small hydrophobic (SH), glycoprotein (G), fusion (F), M2(ORF1), M2(ORF2), and L, substantially as described in Mink et al., Virology 185: 615-624 (1991), Stec et al., Virology 183: 273-287 (1991), and Connors et al., Virol. 208:478-484 (1995), incorporated herein by reference. For purposes of the present invention the genome or antigenome of the recombinant RSV of the invention need only contain those genes or portions thereof necessary to render the viral or subviral particles encoded thereby infectious. Further, the genes or portions thereof may be provided by more than one polynucleotide molecule, i.e., a gene may be provided by complementation or the like from a separate nucleotide molecule.

By recombinant RSV is meant a RSV or RSV-like viral or subviral particle derived directly or indirectly from a recombinant expression system or propagated from virus or subviral particles produced therefrom. The recombinant

expression system will employ a recombinant expression vector which comprises an operably linked transcriptional unit comprising an assembly of at least a genetic element or elements having a regulatory role in RSV gene expression, for example, a promoter, a structural or coding sequence which is transcribed into RSV RNA, and appropriate transcription initiation and termination sequences.

To produce infectious RSV from cDNA-expressed genome or antigenome, the genome or antigenome is coexpressed with those RSV proteins necessary to (i) produce a nucleocapsid capable of RNA replication, and (ii) render progeny nucleocapsids competent for both RNA replication and transcription. Transcription by the genome nucleocapsid provides the other RSV proteins and initiates a productive infection. Alternatively, additional RSV proteins needed for a productive infection can be supplied by coexpression.

An RSV antigenome may be constructed for use in the present invention by, e.g., assembling cloned cDNA segments, representing in aggregate the complete antigenome, by polymerase chain reaction (PCR; described in, e.g., U.S. Patent Nos. 4,683,195 and 4,683,202, and *PCR Protocols: A Guide to Methods and Applications*, Innis et al., eds., Academic Press, San Diego (1990), incorporated herein by reference) of reverse-transcribed copies of RSV mRNA or genome RNA. For example, cDNAs containing the lefthand end of the antigenome, spanning from an appropriate promoter (e.g., T7 RNA polymerase promoter) and the leader region complement to the SH gene, are assembled in an appropriate expression vector, such as a plasmid (e.g., pBR322) or various available cosmid, phage, or DNA virus vectors. The vector may be modified by mutagenesis and/or insertion of synthetic polylinker containing unique restriction sites designed to facilitate assembly. For example, a plasmid vector described herein was derived from pBR322 by replacement of the PstI-EcoRI fragment with a synthetic DNA containing convenient restriction enzyme sites. pBR322 stabilized nucleotides 3716-3732 of the RSV sequence, which otherwise sustained nucleotide

deletions or insertions, and propagation of the plasmid was in bacterial strain DH10B to avoid an artifactual duplication and insertion which otherwise occurred in the vicinity of nt4499. For ease of preparation the G, F and M2 genes can be assembled in a separate vector, as are the L and trailer sequences. The righthand end (e.g., L and trailer sequences) of the antigenome plasmid may contain additional sequences as desired, such as a flanking ribozyme and tandem T7 transcriptional terminators. The ribozyme can be hammerhead type (e.g., Grosfeld et al., J. Virol. 69:5677-5686 (1995)), which would yield a 3' end containing a single nonviral nucleotide, or can any of the other suitable ribozymes such as that of hepatitis delta virus (Perrotta et al., Nature 350:434-436 (1991)) which would yield a 3' end free of non-RSV nucleotides. A middle segment (e.g., G-to-M2 piece) is inserted into an appropriate restriction site of the leader-to-SH plasmid, which in turn is the recipient for the L-trailer-ribozyme-terminator piece, yielding a complete antigenome. In an illustrative example shown in Fig. 1A, the leader end was constructed to abut the promoter for T7 RNA polymerase which included three transcribed G residues for optimal activity; transcription donates these three nonviral G's to the 5' end of the antigenome. These three nonviral G residues can be omitted to yield a 5' end free of nonviral nucleotides. To generate a nearly-correct 3' end, the trailer end was constructed to be adjacent to a hammerhead ribozyme, which upon cleavage would donate a single 3'-phosphorylated U residue to the 3' end of the encoded RNA.

A variety of nucleotide insertions and deletions can be made in the RSV genome or antigenome. The nucleotide length of the genome of wild type human RSV (15,222 nucleotides) is a multiple of six, and members of the Paramyxovirus and Morbillivirus genera typically abide by a "rule of six," i.e., genomes (or minigenomes) replicate efficiently only when their nucleotide length is a multiple of six (thought to be a requirement for precise spacing of nucleotide residues relative to encapsidating NP protein).

Alteration of RSV genome length by single residue increments had no effect on the efficiency of replication, and sequence analysis of several different minigenome mutants following passage showed that the length differences were maintained without compensatory changes. Thus, RSV lacks the strict requirement of genome length being a multiple of six, and nucleotide insertions and deletions can be made in the RSV genome or antigenome without defeating replication of the recombinant RSV of the present invention.

Alternative means to construct cDNA encoding the genome or antigenome include by reverse transcription-PCR using improved PCR conditions (e.g., as described in Cheng et al., Proc. Natl. Acad. Sci. USA 91:5695-5699 (1994)), incorporated herein by reference) to reduce the number of subunit cDNA components to as few as one or two pieces. In other embodiments different promoters can be used (e.g., T3, SP6) or different ribozymes (e.g., that of hepatitis delta virus. Different DNA vectors (e.g., cosmids) can be used for propagation to better accommodate the larger size genome or antigenome.

By virtue of the present invention a variety of alterations in the RSV genome or antigenome for incorporation into infectious recombinant RSV are made possible. For example, foreign genes may be inserted, the order of genes changed, gene overlap removed, the RSV genome promoter replaced with its antigenome counterpart, portions of genes removed (e.g., the cytoplasmic tails of glycoprotein genes), and even entire genes deleted. Modifications in the sequence can be made to facilitate manipulations, such as the insertion of unique restriction sites in various intergenic regions (e.g., a unique *Stu*I site between the G and F genes) or elsewhere. Nontranslated gene sequences can be removed to increase capacity for inserting foreign sequences.

The infectious RSV produced from cDNA-expressed genome or antigenome can be any of the RSV or RSV-like strains, e.g., human, bovine, murine, etc., or of any pneumovirus, e.g., pneumonia virus of mice or turkey

rhinotracheitis virus. To engender a protective immune response, the RSV strain may be one which is endogenous to the subject being immunized, such as human RSV being used to immunize humans. The genome or antigenome can be modified, however, to express RSV genes from different types. Thus, infectious RSV intended for administration to humans may be human RSV that has been modified to contain genes from a bovine or murine RSV type such as for the purpose of attenuation, or a bovine RSV may be modified to contain genes that encode epitopes or proteins that elicit protection against human RSV infection, e.g., the human RSV glycoprotein genes can be substituted for the bovine glycoprotein genes such that the bovine RSV, which has a restricted ability to replicate in a human host, elicits a protective immune response in humans against human RSV strains.

The N, P and L proteins, necessary for RNA replication, require an RNA polymerase elongation factor such as the M2(ORF1) protein for processive transcription. Thus M2(ORF1) or a substantially equivalent transcription elongation factor for negative strand RNA viruses is required for the production of infectious RSV and is a necessary component of functional nucleocapsids during productive infection. The need for the M2(ORF1) protein is consistent with its role as a transcription elongation factor. The need for expression of the RNA polymerase elongation factor protein for negative strand RNA viruses is a feature of the present invention. M2(ORF1) can be supplied by expression of the complete M2-gene, although in this form the second ORF2 may also be expressed and have an inhibitory effect on RNA replication. Therefore, for production of infectious virus using the complete M2 gene the activities of the two ORFs should be balanced to permit sufficient expression of M(ORF1) to provide transcription elongation activity yet not so much of M(ORF2) to inhibit RNA replication. Alternatively, the ORF1 protein is provided from a cDNA engineered to lack ORF2 or which encodes a defective ORF2. Efficiency of virus production may also be improved by co-expression of additional

viral protein genes, such as those encoding envelope constituents (i.e., SH, M, G, F proteins).

Isolated polynucleotides (e.g., cDNA) encoding the genome or antigenome and, separately, the N, P, L and M2(ORF1) proteins, are inserted by transfection, electroporation, mechanical insertion, transduction or the like, into cells which are capable of supporting a productive RSV infection, e.g., HEP-2, FRhL-DBS2, MRC, and Vero cells. Transfection of isolated polynucleotide sequences may be introduced into cultured cells by, for example, calcium phosphate-mediated transfection (Wigler et al., Cell 14: 725 (1978); Corsaro and Pearson, Somatic Cell Genetics 7: 603 (1981); Graham and Van der Eb, Virology 52: 456 (1973)), electroporation (Neumann et al., EMBO J. 1: 841-845 (1982)), DEAE-dextran mediated transfection (Ausubel et al., (ed.) Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY (1987), incorporated herein by reference), cationic lipid-mediated transfection (Hawley-Nelson et al., Focus 15: 73-79 (1993)) or a commercially available transfection reagent, e.g., LipofectACE® (Life Technologies). The N, P, L and M2(ORF1) proteins are encoded by one or more expression vectors which can be the same or separate from that which encodes the genome or antigenome, and various combinations thereof. Additional proteins may be included as desired, encoded on a its own vector or on a vector encoding a N, P, L, or M2(ORF1) protein or the complete genome or antigenome. Expression of the genome or antigenome and proteins from transfected plasmids can be achieved, for example, by each cDNA being under the control of a promoter for T7 RNA polymerase, which in turn is supplied by infection, transfection or transduction with an expression system for the T7 RNA polymerase, e.g., a vaccinia virus MVA strain recombinant which expresses the T7 RNA polymerase (Wyatt et al., Virology, 210: 202-205 (1995), incorporated herein by reference). The viral proteins, and/or T7 RNA polymerase, can also be provided from transformed mammalian cells, or by transfection of preformed mRNA or protein.

Alternatively, synthesis of antigenome or genome together with the above-mentioned viral proteins can be done in vitro (cell-free) in a combined transcription-translation reaction, followed by transfection into cells. Or, antigenome or genome RNA can be synthesized in vitro and transfected into cells expressing RSV proteins.

Having the infectious clone of the invention permits the alteration of the RSV genome (or antigenome) by introducing defined mutations. By "infectious clone" is meant cDNA or its product, synthetic or otherwise, which can be transcribed into genomic or antigenomic RNA capable of serving as template to produce the genome of infectious virus or subviral particle. Defined mutations can be introduced by conventional techniques (e.g., site-directed mutagenesis) into a cDNA copy of the genome or antigenome. The use of antigenome cDNA subfragments to assemble a complete antigenome cDNA as described herein has the advantage that each region can be manipulated separately (smaller cDNAs are easier to manipulate than large ones) and then readily assembled into a complete cDNA. Thus, the complete antigenome or genome cDNA, or any subfragment thereof, can be used as template for oligonucleotide-directed mutagenesis. This can be through the intermediate of a single-stranded phagemid form, such as using the Muta-gen kit of Bio-Rad, or a method using the double-stranded plasmid directly as template such as the Chameleon mutagenesis kit of Stratagene, or by the polymerase chain reaction employing either an oligonucleotide primer or template which contains the mutation(s) of interest. A mutated subfragment can then be assembled into the complete antigenome or genome cDNA. A variety of other mutagenesis techniques are known and available for use in producing the mutations of interest in the RSV antigenome or genome cDNA. Mutations can vary from single nucleotide changes to replacement of large cDNA pieces containing one or more genes or genome regions.

Thus, in one illustrative embodiment mutations are introduced by using the Muta-gene phagemid in vitro

mutagenesis kit available from Bio-Rad Laboratories, Richmond, CA. In brief, cDNA encoding an RSV genome or antigenome is cloned into the plasmid pTZ18U, and used to transform DH5 α F' cells (Life Technologies Inc., Gaithersburg, MD). Phagemid preparations are prepared as recommended by the manufacturer. 5 Oligonucleotides are designed for mutagenesis by introduction of an altered nucleotide at the desired position of the genome or antigenome. The plasmid containing the genetically altered genome or antigenome is then amplified.

10 The ability to introduce defined mutations into infectious RSV has many applications, including the analyses of RSV molecular biology and pathogenesis. For example, the functions of the RSV proteins, including the NS1, NS2, SH, M2(ORF1) and M2(ORF2) proteins, can be investigated by 15 introducing mutations which ablate or reduce their level of expression, or which yield mutant protein.

As another example, the sequence at the cleavage site of the F protein, or the putative attachment domain of the G protein, can be modified to evaluate effects on growth 20 in tissue culture and infection and pathogenesis in experimental animals.

The roles of various genome RNA structural features, such as promoters, intergenic regions, gene overlap, and transcription signals, can be evaluated using the methods 25 and compositions of the present invention. Evaluation of trans-acting proteins and cis-acting RNA sequences using the complete antigenome cDNA can be conducted in parallel using RSV minigenomes (e.g., Grosfeld et al., J. Virol. 69: 5677-5686 (1995), incorporated herein by reference), whose 30 helper-dependent status is useful in the characterization of those mutants which are too inhibitory to be recovered in replication-independent infectious virus.

A number of attenuated RSV strains as candidate vaccines for intranasal administration have been developed 35 using multiple rounds of chemical mutagenesis to introduce multiple mutations into a virus which had already been attenuated during cold-passages (e.g., Connors et al., Virology

208: 478-484 (1995); Crowe et al., Vaccine 12: 691-699 (1994);
and Crowe et al., Vaccine 12: 783-790 (1994), incorporated
herein by reference). Evaluation in rodents, chimpanzees,
adults and infants indicate that certain of these candidate
5 vaccine strains are relatively stable genetically, are highly
immunogenic, and may be satisfactorily attenuated. Nucleotide
sequence analysis of some of these attenuated viruses
indicates that each level of increased attenuation is
associated with two or more new nucleotide and amino acid
10 substitutions (Connors et al., supra). The present invention
provides the ability to distinguish between silent incidental
mutations versus those responsible for phenotype differences
by introducing the mutations, separately and in various
combinations, into the genome or antigenome of infectious
15 wild-type RSV. This process identifies mutations responsible
for phenotypes such as attenuation, temperature sensitivity,
cold-adaptation, small plaque size, host range restriction,
etc. Mutations from this menu can then be introduced in
various combinations to calibrate a vaccine virus to an
20 appropriate level of attenuation, etc., as desired. Moreover,
the present invention provides the ability to combine
mutations from different strains of virus into one strain.

The present invention also provides new methods of
attenuation. For example, individual internal genes of human
25 RSV can be replaced with their bovine, murine or other RSV
counterpart. This may include part or all of one or more of
the NS1, NS2, N, P, M, SH, M2(ORF1), M2(ORF2) and L genes, or
non-immunogenic parts of the G and F genes. Reciprocally,
means are provided to generate a live attenuated bovine RSV by
30 inserting human attenuating genes into a bovine RSV genome or
antigenome background. Human RSV bearing bovine RSV
glycoproteins provides a host range restriction favorable for
human vaccine preparations. Bovine RSV sequences which can be
used in the present invention are described in, e.g., Pastey
35 et al., J. Gen. Virol. 76:193-197 (1993); Pastey et al., Virus
Res. 29:195-202 (1993); Zamora et al., J. Gen. Virol. 73:737-
741 (1992); Mallipeddi et al., J. Gen. Virol. 74:2001-2004

(1993); Mallipeddi et al., J. Gen. Virol. 73:2441-2444 (1992); and Zamora et al., Virus Res. 24:115-121 (1992), each of which is incorporated herein by reference.

The invention also provides the ability to analyze other types of attenuating mutation and to incorporate them into infectious RSV for vaccine or other uses. For example, a tissue culture-adapted nonpathogenic strain of pneumonia virus of mice (the murine counterpart of RSV) lacks a cytoplasmic tail of the G protein (Randhawa et al., Virology 207: 240-245 (1995)). By analogy, the cytoplasmic and transmembrane domains of each of the RSV glycoproteins, F, G and SH, can be deleted or modified to achieve attenuation.

Other mutations for use in infectious RSV of the present invention include mutations in cis-acting signals identified during mutational analysis of RSV minigenomes. For example, insertional and deletional analysis of the leader and trailer and flanking sequences identified viral promoters and transcription signals and provided a series of mutations associated with varying degrees of reduction of RNA replication or transcription. Saturation mutagenesis (whereby each position in turn is modified to each of the nucleotide alternatives) of these cis-acting signals also has identified many mutations which reduced (or in one case increased) RNA replication or transcription. Any of these mutations can be inserted into the complete antigenome or genome as described herein. Other mutations involve replacement of the 3' end of genome with its counterpart from antigenome, which is associated with changes in RNA replication and transcription. In addition, the intergenic regions (Collins et al., Proc. Natl. Acad. Sci. USA 83:4594-4598 (1986), incorporated herein by reference) can be shortened or lengthened or changed in sequence content, and the naturally-occurring gene overlap (Collins et al., Proc. Natl. Acad. Sci. USA 84:5134-5138 (1987), incorporated herein by reference) can be removed or changed to a different intergenic region by the methods described herein.

In another embodiment, RSV useful in a vaccine formulation can be conveniently modified to accommodate antigenic drift in circulating virus. Typically the modification will be in the G and/or F proteins. The entire G or F gene, or the segment(s) encoding particular immunogenic regions thereof, is incorporated into the RSV genome or antigenome cDNA by replacement of the corresponding region in the infectious clone or by adding one or more copies of the gene such that several antigenic forms are represented.

Progeny virus produced from the modified RSV cDNA are then used in vaccination protocols against the emerging strains. Further, inclusion of the G protein gene of RSV subgroup B would broaden the response to cover a wider spectrum of the relatively diverse subgroup A and B strains present in the human population.

An infectious RSV clone of the invention can also be engineered to enhance its immunogenicity and induce a level of protection greater than that provided by natural infection, or vice versa, to identify and ablate epitopes associated with undesirable immunopathologic reactions. Enhanced immunogenicity of the vaccines produced by the present invention addresses one of the greatest obstacles to controlling RSV, namely the incomplete nature of immunity induced by natural infection. An additional gene may be inserted into or proximate to the RSV genome or antigenome which is under the control of an independent set of transcription signals. Genes of interest include those encoding cytokines (e.g., IL-2 through IL-15, especially IL-3, IL-6 and IL-7, etc.), gamma-interferon, and proteins rich in T helper cell epitopes. The additional protein can be expressed either as a separate protein or as a chimera engineered from a second copy of one of the RSV proteins, such as SH. This provides the ability to modify and improve the immune response against RSV both quantitatively and qualitatively.

For vaccine use, virus produced according to the present invention can be used directly in vaccine formulations, or lyophilized, as desired, using lyophilization

protocols well known to the artisan. Lyophilized virus will typically be maintained at about 4°C. When ready for use the lyophilized virus is reconstituted in a stabilizing solution, e.g., saline or comprising SPG, Mg⁺⁺ and HEPES, with or without adjuvant, as further described below.

Thus RSV vaccines of the invention contain as an active ingredient an immunogenetically effective amount of RSV produced as described herein. The modified virus may be introduced into a host with a physiologically acceptable carrier and/or adjuvant. Useful carriers are well known in the art, and include, e.g., water, buffered water, 0.4% saline, 0.3% glycine, hyaluronic acid and the like. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration, as mentioned above. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, and the like. Acceptable adjuvants include incomplete Freund's adjuvant, aluminum phosphate, aluminum hydroxide, or alum, which are materials well known in the art.

Upon immunization with a RSV composition as described herein, via aerosol, droplet, oral, topical or other route, the immune system of the host responds to the vaccine by producing antibodies specific for RSV virus proteins, e.g., F and G glycoproteins. As a result of the vaccination the host becomes at least partially or completely immune to RSV infection, or resistant to developing moderate or severe RSV infection, particularly of the lower respiratory tract.

The host to which the vaccine are administered can be any mammal which is susceptible to infection by RSV or a closely related virus and which host is capable of generating a protective immune response to the antigens of the

vaccinizing strain. Thus, suitable hosts include humans, non-human primates, bovine, equine, swine, ovine, caprine, lagamorph, rodents, etc. Accordingly, the invention provides methods for creating vaccines for a variety of human and veterinary uses.

The vaccine compositions containing the RSV of the invention are administered to a host susceptible to or otherwise at risk of RSV infection to enhance the host's own immune response capabilities. Such an amount is defined to be a "immunogenically effective dose." In this use, the precise amounts again depend on the host's state of health and weight, the mode of administration, the nature of the formulation, etc., but generally range from about 10^3 to about 10^6 plaque forming units (PFU) or more of virus per host, more commonly from about 10^4 to 10^5 PFU virus per host. In any event, the vaccine formulations should provide a quantity of modified RSV of the invention sufficient to effectively protect the host patient against serious or life-threatening RSV infection.

The RSV produced in accordance with the present invention can be combined with viruses of the other subgroup or strains to achieve protection against multiple RSV subgroups or strains, or protective epitopes of these strains can be engineered into one virus as described herein. Typically the different viruses will be in admixture and administered simultaneously, but may also be administered separately. For example, as the F glycoproteins of the two RSV subgroups differ by only about 11% in amino acid sequence, this similarity is the basis for a cross-protective immune response as observed in animals immunized with RSV or F antigen and challenged with a heterologous strain. Thus, immunization with one strain may protect against different strains of the same or different subgroup.

In some instances it may be desirable to combine the RSV vaccines of the invention with vaccines which induce protective responses to other agents, particularly other childhood viruses. For example, the RSV vaccine of the present invention can be administered simultaneously with

parainfluenza virus vaccine, such as described in Clements et al., J. Clin. Microbiol. 29:1175-1182 (1991), which is incorporated herein by reference. In another aspect of the invention the RSV can be employed as a vector for protective antigens of other respiratory tract pathogens, such as parainfluenza, by incorporating the sequences encoding those protective antigens into the RSV genome or antigenome which is used to produce infectious RSV as described herein.

Single or multiple administrations of the vaccine compositions of the invention can be carried out. In neonates and infants, multiple administration may be required to elicit sufficient levels of immunity. Administration should begin within the first month of life, and at intervals throughout childhood, such as at two months, six months, one year and two years, as necessary to maintain sufficient levels of protection against native (wild-type) RSV infection. Similarly, adults who are particularly susceptible to repeated or serious RSV infection, such as, for example, health care workers, day care workers, family members of young children, the elderly, individuals with compromised cardiopulmonary function, may require multiple immunizations to establish and/or maintain protective immune responses. Levels of induced immunity can be monitored by measuring amounts of neutralizing secretory and serum antibodies, and dosages adjusted or vaccinations repeated as necessary to maintain desired levels of protection. Further, different vaccine viruses may be advantageous for different recipient groups. For example, an engineered RSV strain expressing an additional protein rich in T cell epitopes may be particularly advantageous for adults rather than for infants.

In yet another aspect of the invention the RSV is employed as a vector for transient gene therapy of the respiratory tract. According to this embodiment the recombinant RSV genome or antigenome incorporates a sequence which is capable of encoding a gene product of interest. The gene product of interest is under control of the same or a different promoter from that which controls RSV expression.

The infectious RSV produced by coexpressing the recombinant RSV genome or antigenome with the N, P, L and M2(ORF1) proteins and containing a sequence encoding the gene product of interest is administered to a patient. Administration is typically by aerosol, nebulizer, or other topical application to the respiratory tract of the patient being treated. Recombinant RSV is administered in an amount sufficient to result in the expression of therapeutic or prophylactic levels of the desired gene product. Examples of representative gene products which are administered in this method include those which encode, for example, those particularly suitable for transient expression, e.g., interleukin-2, interleukin-4, gamma-interferon, GM-CSF, G-CSF, erythropoietin, and other cytokines, glucocerebrosidase, phenylalanine hydroxylase, cystic fibrosis transmembrane conductance regulator (CFTR), hypoxanthine-guanine phosphoribosyl transferase, cytotoxins, tumor suppressor genes, antisense RNAs, and vaccine antigens.

The following examples are provided by way of illustration, not limitation.

EXAMPLE I

Construction of cDNA Encoding RSV Antigenome

A cDNA clone encoding the antigenome of RSV strain A2 was constructed, as shown in Fig. 1A. The cDNA was synthesized in segments by reverse transcription (RT) and polymerase chain reaction (PCR) using synthetic oligonucleotides as primers and intracellular RSV mRNA or genome RNA isolated from purified virions as template. The final cDNA was flanked on the leader end by the promoter for T7 RNA polymerase, which included three transcribed G residues for optimal activity; transcription would result in the donation of these three nonviral G's to the 5' end of the antigenome. To generate a nearly-correct 3' end, the cDNA trailer end was constructed to be adjacent to a previously-described hammerhead ribozyme, which upon cleavage would donate a single 3'-phosphorylated U residue to the 3'

end of the encoded RNA (Grosfeld et al., J. Virol. 69:5677-5686, incorporated herein by reference). The ribozyme sequence was followed by a tandem pair of terminators of T7 RNA polymerase. (The addition of three 5' G residues and one 3' U residue to a cDNA-encoded RSV minigenome containing the chloramphenicol acetyl transferase (CAT) reporter gene had no effect on the expression of CAT when complemented by RSV.)

Fig. 1A shows the structures of the cDNA and the encoded antigenome RNA (not to scale). The diagram of the antigenome (at top) includes the following features: the 5'-terminal nonviral G triplet contributed by the T7 promoter, the four sequence markers at positions 1099 (which adds one nt to the length), 1139, 5611, and 7559, the ribozyme and tandem T7 terminators, and the single nonviral 3'-phosphorylated U residue contributed to the 3' end by ribozyme cleavage (the site of cleavage is indicated with an arrow) (13).

Cloned cDNA segments (Fig. 1A, middle) representing in aggregate the complete antigenome were constructed by RT-PCR of RSV mRNA or genome RNA. cDNAs containing the lefthand end of the antigenome, spanning from the T7 promoter and leader region complement to the SH gene, were assembled in a version of pBR322 (Fig. 1A, bottom) in which the naturally-occurring BamHI site had been ablated by mutagenesis and the PstI-EcoRI fragment replaced with a synthetic polylinker containing unique restriction sites (including BstBI, BstXI, PacI, BamHI, MluI) designed to facilitate assembly. The box in Fig. 1A shows the removal of the BamHI site. The naturally occurring BamHI-SalI fragment (the BamHI site is shown in the top line in positive sense, underlined) was replaced with a PCR-generated BglII-SalI fragment (the BglII site is shown in the bottom line, underlined; its 4-nt sticky end [italics] is compatible with that of BamHI). This resulted in a single nt change (middle line, underlined) which was silent at the amino acid level. These modifications to the vector facilitated construction of the cDNA by rendering unique a BamHI site in the antigenome cDNA.

The G, F and M2 genes were assembled in a separate plasmid, as were the L, trailer and flanking ribozyme and tandem T7 transcription terminators. The G-to-M2 piece was then inserted into the PacI-BamHI window of the leader-to-SH plasmid. This in turn was the recipient for the L-trailer-ribozyme-terminator piece inserted into the BamHI to MluI, yielding the complete antigenome.

Four restriction site markers (Fig. 1B) were introduced into the antigenome cDNA by incorporating the changes into oligonucleotide primers used in RT-PCR. This was done to facilitate assembly, provide a means to identify recombinant virus, and illustrate the ability to introduce changes into infectious RSV. Three sites were in intergenic regions and the fourth in a nontranslated gene region, and they involved a total of five nt substitutions and a single nt insertion. This increased the length of the encoded antigenome by one nt from that of wild type to a total of 15,223 nt (SEQ ID NO:1, which depicts the 5' to 3' positive-sense sequence whereas the genome itself is negative-sense).

The sequence markers were inserted into the cDNA-encoded antigenome RNA as shown in Fig. 1B. Sequences are positive sense and numbered relative to the first nt of the leader region complement as 1; identities between strains A2 and 18537 (Johnson and Collins, J. Gen Virol. 69:2901-2906 (1988), incorporated herein by reference), representing subgroups A and B, respectively, are indicated with dots; sequences representing restriction sites in the cDNA are underlined; gene-start (GS) and gene-end (GE) transcription signals are boxed; the initiation codon of the N translational open reading frame at position 1141 is italicized, and the sequence markers are shown underneath each sequence. In the top sequence, a single C residue was inserted at position 1099 to create an AflII site in the NSII-N intergenic region, and the AG at positions 1139 and 1140 immediately upstream of the N translational open reading frame were replaced with CC to create a new NcoI site. In the middle sequence, substitution of G and U at positions 5612 and 5616, respectively, created a

new StuI site in the G-F intergenic region. And, in the bottom sequence of Fig. 1B, a C replacement at position 7561 created a new SphI site in the F-M2 intergenic region.

All cDNAs were sequenced in their entirety, in most instances from several independent cDNAs, prior to assembly. The plasmids encoding individual RSV proteins are described in Grosfeld et al., J. Virol. 69:5677-5686 (1995) and Collins et al., Proc. Natl. Acad. Sci. USA (1995), each of which is incorporated herein by reference.

EXAMPLE II

Transfection and Recovery of Recombinant RSV

The strategy for producing infectious RSV from cDNA-expressed antigenome involved its coexpression with those RSV proteins which are sufficient to (i) produce an antigenome nucleocapsid capable of RNA replication, and (ii) render the progeny genome nucleocapsid competent for both RNA replication and transcription. Transcription by the genome nucleocapsid provides all of the other RSV proteins and initiates a productive infection.

Plasmid-borne cDNA encoding the antigenome was transfected, together with plasmids encoding proteins N, P, L and M2(ORF1), into HEp-2 cells which had been infected with a recently-described vaccinia virus MVA strain recombinant which expresses the T7 RNA polymerase (Wyatt et al., Virol. 210:202-205 (1995), incorporated herein by reference). The MVA strain is a host range mutant which grows permissively in avian cells whereas in mammalian cells there is a block at a late stage in virion maturation that greatly reduces the production of infectious virus. In HEp-2 cells, the MVA recombinant was similar to the more commonly-used WR-based recombinant (Fuerst et al., Proc. Natl. Acad. Sci. USA 83: 8122-8126 (1986)) with regard to the level of expression of T7 polymerase and cytopathogenicity, but the level of progeny produced was sufficiently low that supernatants could be passaged to fresh

cells with minimal cytopathogenicity. This should facilitate the recovery of any recombinant RSV which might be produced in transfected, vaccinia virus-infected cells.

Transfection and recovery of recombinant RSV was performed as follows. Monolayer cultures of HEp-2 cells received, per single well of a six-well dish, one ml of infection-transfection medium prepared by mixing five plasmids in a final volume of 0.1 ml Opti-MEM (Life Technologies) medium, namely 0.4 μ g each of antigenome, N and P plasmids, and 0.1 μ g each of L and M2(ORF1) plasmids. This was combined with 0.1 ml of Opti-MEM containing 12 μ l LipofectACE (Life Technologies). After 15 min incubation at room temperature, this was combined with 0.8 ml of OptiMEM containing 2% heat-inactivated fetal calf serum and 1.5×10^6 pfu of strain MVA vaccinia virus recombinant encoding T7 RNA polymerase (Wyatt et al., supra). This was added to the cells and replaced one day later by Opti-MEM containing 2% serum. Cultures were incubated at 32°C and harvested on day three. Incubation at 32°C was used because it was found that the MVA virus is slightly temperature sensitive and is much more efficient at this lower temperature.

Three days post-transfection clarified culture supernatants were passaged onto fresh HEp-2 cells and overlaid with methyl cellulose (for subsequent antibody staining) or agarose (for plaque isolation). After incubation for five days under methyl cellulose, the cells were fixed and stained by an indirect horseradish peroxidase method using a mixture of three murine monoclonal antibodies to the RSV F protein followed by an anti-mouse antibody linked to horseradish peroxidase, following the general procedure of Murphy et al., Vaccine 8: 497-502 (1990).

Numerous RSV-like plaques were detected against a background of cytopathogenicity that presumably was due to a low level of MVA-T7 recombinant virus. The plaques contained an abundant amount of the RSV F protein, as indicated by brown-black coloration, and displayed cytopathic effects characteristic of RSV, notably syncytium formation.

The RSV-like plaques were picked from plates which had been prepared in parallel but incubated under agarose and stained with neutral red. They were propagated and compared to a laboratory strain of RSV strain A2 by plaque assay and antibody staining. The plaques derived from the transfected cultures closely resembled those of the laboratory strain. One difference was that the plaques derived from the transfected cultures appeared to be slightly smaller than those from the laboratory strain, with centers which were less well cleared. The recombinant virus may differ phenotypically from this particular wild type isolate, possibly being slightly more restricted in cell-to-cell spread and exhibiting a reduced rate of cell killing. With regard to the propagation of released virus, the yields of the recombinant versus laboratory virus in HEP-2 cells were essentially identical at 32° or 37°C. In preliminary studies, the recombinant and laboratory viruses were indistinguishable with regard to the accumulation of intracellular RSV mRNAs and proteins.

Plaque-purified, thrice-passaged recombinant RSV was analyzed in parallel with laboratory virus by RT-PCR using three primer pairs flanking the four inserted markers. Three independent plaque-purified recombinant RSV isolates were propagated in parallel with an uninfected control culture. Clarified medium supernatants were treated with polyethylene glycol and high salt (Zoller and Smith, DNA 3:479-488 (1984)) to precipitate virus and RNA was extracted from the pellets with TrizolTM (Life Technologies). These RNAs, in parallel with additional controls of no added RNA or 0.1 µg of RNA from a laboratory isolate of strain A2, were treated with DNase, repurified, annealed each with 50 ng of random hexamers and incubated under standard RT conditions (40 µl reactions) with or without reverse transcriptase (Connors et al., Virology 208: 478-484 (1995)). Aliquots of each reaction were subjected to PCR (35 cycles of 94°C for 45s, 37°C for 30s, 72°C for 1 min) using three different pairs of synthetic deoxyoligonucleotide primers. Primer pair (A): positive-sense, positions 925-942

and negative-sense, positions 1421-1440, yielding a predicted product of 516 bp (517 bp in the case of the recombinant viruses) that included the AflII and NcoI sites inserted at, respectively, the junction of the NS2 and N genes and in the N gene. Primer pair (B): positive-sense, positions 5412-5429 and negative-sense, 5930-5949, yielding a predicted product of 538 bp spanning the StuI site inserted at the junction between the G and F genes. Primer pair (C): positive-sense, 7280-7297 and negative-sense, 7690-7707, yielding a 428 bp fragment spanning the SphI site inserted at the junction between the F and M2 genes. PCR products were analyzed by electrophoresis on neutral gels containing 1% agarose and 2% low-melting agarose in parallel with HaeIII-digested X174 DNA molecular length markers and visualized by staining with ethidium bromide. PCR products of the expected sizes were produced. The production of each was dependent on the RT step, indicating that each was derived from RNA rather than contaminating cDNA.

PCR products were analyzed by digestion with restriction enzymes. Digestion of products of primer pair A with AflII or NcoI yielded fragments corresponding to the predicted 177 and 340 bp (AflII) or 217 and 300 bp (NcoI). Digestion of products of primer pair B with StuI yielded fragments comparable to the predicted 201 and 337 bp. Digestion of products from reactions with primer pair C with SphI yielded products corresponding to the predicted 147 and 281 bp. The digests were analyzed by gel electrophoresis as above. The presence of residual undigested PCR product with AflII was due to incomplete digestion, as was confirmed by redigestion. Thus, the restriction enzyme digestion showed that the PCR products representing recombinant virus contained the expected restriction site markers while those representing the laboratory strain did not. Nucleotide sequence analysis of cloned PCR product confirmed the sequences spanning the restriction site markers.

As shown in Table 1, the efficiency of RSV production when complemented by N, P, L and M2(ORF1) was

relatively high, ranging in three experiments from an average of 9.9 to 94.8 plaques per 0.4 μ g of input antigenome cDNA and 1.5 X 10⁶ cells. Since these plaques were derived from passage, the number of infected cells present in each well of the original transfection was not known. Nearly every transfected well (54 of 56 in Table 1) produced virus. Since the yield of released RSV per infected cell typically is very low (~10 pfu) even under ideal conditions, and since many wells yielded many times this amount (up to 169 plaques in Table 1), it is likely that several RSV producing cells were present in many of the wells of transfected cells.

RSV was not recovered if any of the plasmids were omitted (e.g., as shown in Table 1). The requirement for M2(ORF1) also could be satisfied with the complete gene, M2(ORF1+2), provided the level of its input cDNA was low (0.016 μ g per 1.5 X 10⁶ cells [Table 1]). At higher levels, the production of virus was greatly reduced, suggesting that an inhibition of minigenome RNA synthesis associated with M2(ORF2) also operates on the complete genome during productive infection.

These results showed that the production of infectious RSV was highly dependent on expression of the M2(ORF1) protein in addition to N, P and L. Furthermore, it showed that the optimal method of expression of M2(ORF1) was from an engineered cDNA in which ORF2 had been deleted, although the complete cDNA containing both ORFs also supported the production of RSV.

Thus, as part of the present invention, transcription by RSV differed from previously-described nonsegmented negative strand RNA viruses in requiring a fourth protein designated here as M2(ORF1), and previously called 22K or M2 (Collins et al., J. Virol. 54:65-71 (1985)). The M2(ORF1) protein was found to be an RNA polymerase elongation factor essential for processive, sequential transcription. This requirement provides the capability, as part of this invention, for introducing specific, predetermined changes into infectious RSV.

Table 1. Production of infectious RSV was dependent on expression of M2 ORF 1.

5	Complementing plasmids (μ g cDNA per 1.5×10^6 cells and antigenome cDNA) 0.4 μ g	Production of infectious RSV #plaques X # wells		
		expt. 1	expt. 2	expt. 3
10	N(0.4) P(0.4) L(0.1)	0 X 24	0 X 12	0 X 12
15	N(0.4) P(0.4) L(0.1) M2 [ORF1+2] (0.016)	0 X 19 ⁵ 1 X 2 2 X 2 3 X 1	0 X 4 3 X 1 5 X 1 6 X 1 9 X 1	9 X 1 10 X 1 14 X 2 22 X 1 28 X 1
20		av. 0.38	10 X 1 13 X 1 34 X 1 51 X 1	32 X 1 49 X 1 70 X 2 166 X 1 169 X 1
25			av. 10.9	av. 48.6
30	N(0.4) P(0.4) L(0.1) M2 [ORF1] (0.1)	0 X 1 1 X 1 2 X 2 3 X 2 4 X 1 5 X 2 6 X 4 7 X 2 9 X 1 10 X 2	11 X 1 12 X 1 13 X 1 21 X 1 24 X 1 26 X 1 30 X 2 33 X 2 42 X 1 73 X 1	0 X 1 2 X 1 4 X 1 5 X 1 8 X 2 10 X 3 19 X 1 20 X 1 23 X 1 23 X 1
35			av. 9.9	147 X 1 148 X 1
40				av. 94.8

* Supernatants from transfected cultures (10^6 cells per well) were passaged onto fresh HEp-2 cells, overlaid with methyl cellulose, and stained with F-specific monoclonal antibodies.

⁵ Read as follows: 19 wells had 0 plaques, 2 wells had 1 plaque each, 2 wells had 2 plaques each, and 1 well had 3 plaques.

EXAMPLE III

Constructing Infectious RSV With Predetermined Mutations
To Confer a Desired Phenotype

5 This Example illustrates the introduction of
specific predetermined mutations into infectious recombinant
RSV using the methods described hereinabove. For ease of
manipulation, the antigenome cDNA was cloned as two separate
pieces in separate plasmids: one piece (the left end)
10 containing the T7 promoter together with nucleotide 1 through
to the BamHI site at nucleotide 8501 (cDNA D50), and the other
(the right end) containing the BamHI site through to
nucleotide 15223, together with the ribozyme and T7
transcription terminators (cDNA D39). D39 was further
15 separated into two pieces and each placed in a separate
phagemid plasmid: one piece (left hand half, cDNA L1) from the
BamHI site to the PmlI site at nucleotide 12255, and the other
(right hand half, cDNA L2) from the PmlI site to the end of
the T7 terminator. The sequence positions assigned to
20 restriction site locations are intended as a descriptive guide
and do not alone precisely define all of the nucleotides
involved.

 Following a general procedure of Kunkel et al. Meth.
Enzymol. 54:367-382 (1987) (incorporated herein by reference),
25 the plasmids were propagated in a dut ung strain of E. coli,
strain CJ236, and single stranded DNA was prepared by
infection with a helper phage, M13KO7. Phosphorylated
synthetic oligonucleotides each containing one or more
nucleotide changes of interest were prepared, annealed to the
30 single stranded template singly or more than one at a time,
and used to direct DNA synthesis by T4 DNA polymerase. The
products were ligated and transformed into a non-dut ung
strain of E. coli, DH5 α or DH10B. Colonies containing mutant
plasmids were identified by restriction enzyme digestion or
35 by sequence analysis. Other methods of mutagenesis can
readily be used.

L1 and L2 described above were modified to contain several combinations of recognition sites for several different restriction enzymes which do not appear or are infrequent in the antigenome plasmid; these sites were introduced using nucleotide substitutions which did not change the amino acid sequence of the encoded L protein. In addition, L1 was modified to contain a mutation believed to confer a ts phenotype. Two versions of L1 were made. In one version, L1 was modified in a single cycle of mutagenesis to contain new Bsu36I (nucleotide 9399) and SnaBI (11848) sites, and a mutation termed 530, yielding cDNA 530L1sites. The 530 mutation had been identified by sequence analysis of the biologically-derived virus cpts530-RSV and involves a single nucleotide change at position 10060 which results in an amino acid change (Phe to Leu) at amino acid 521 in the L protein. In a second version, L1 was modified in a single cycle to contain the Bsu36I and SnaBI sites, resulting in cDNA L1 sites. L2 was modified in one cycle of mutagenesis to contain the new sites PmeI (13342), RsrII (14083) and SnaBI (14477). In a second cycle, the site BstEII (14318) was added and a naturally-occurring recognition site for SnaBI (6956) was removed. This yielded L2 sites.

Three complete antigenome cDNAs were made by introducing selected L1 and L2 mutant cDNAs or fragments thereof into D39 and combining this with D50. Antigenome cDNA "D53sites" contains L1 sites and L2 sites. cDNA "530D53" contains the BamHI-SpeI (10149) fragment of 530L1sites (which contains the Bsu36I and 530 mutations). cDNA "530D53sites" contained 530L1 sites and L2 sites (Table II). Recombinant virus was recovered from each of the three complete mutant antigenome cDNAs using the methodology of this invention and were passaged at least twice and analyzed directly or following plaque purification and amplification. The presence of mutations was confirmed by RT-PCR of viral RNA followed by analysis by restriction enzyme digestion or nucleotide sequencing, or both.

The engineered viruses were evaluated for their ability to form plaques in HEP-2 cells at 32°C, 39°C and 40°C in parallel with two nonrecombinant biologically-derived viruses, HEK, a wild type strain A2 virus, and cpts530 RSV, the virus from which the 530 mutation was identified by sequence analysis (Table II). This comparison showed that all three engineered viruses formed plaques at 32°C, and showed that the titers of the various virus preparations were within two log₁₀ units of each other, which is within the range of experimental variation typically seen among independent preparations of RSV. The recombinant viruses containing the 530 mutation were greatly impaired in ability to form plaques at 39°C or 40°C, comparable to cpts530 RSV. The presence of additional restriction sites in 530D53sites versus 530D53 had no discernable effect on the ts phenotype. D53sites virus, which contained silent restriction site markers but lacked the 530 mutation, retained the ability to form plaques at the higher temperatures, comparable to wild type. This not only provided positive identification that the 530 mutation is involved in the ts phenotype of cpts530 RSV, but also showed that point mutations can be introduced systematically into recombinant RSV according to the present invention. In these cases, the resulting phenotypes of the engineered viruses were fully consistent with the parental strain and provided direct confirmation and reconstitution of an attenuation mutation.

Table II

Characterization of the ts phenotype of biologically-derived RSV versus RSV recovered from cDNA clones

virus	Efficiency of plaque formation (log ₁₀) at the indicated temperature		
	32°C	39°C	40°C
<u>Biologically-derived viruses</u>			
HEK ^a	8.7	8.6	8.5
cpts530RSV ^b	6.8	<0.7	<0.7
<u>cDNA-derived viruses</u>			
D53 sites ^c	6.9	6.9	6.7
530D53 ^d	7.9	<0.7	<0.7
530D53 sites ^e	7.6	<0.7	<0.7

^a = Wild type RSV A2.

^b = ts virus.

^c = Contains six new restriction sites and lacks one naturally-occurring site.

^d = Contains one new restriction site and the 530 mutation.

^e = Contains six new restriction sites and the 530 mutation and lacks one naturally-occurring site.

EXAMPLE IV

Recovery of Infectious Respiratory Syncytial Virus Expressing An Additional, Foreign Gene

The methods described above were used to construct recombinant RSV containing an additional gene, encoding chloramphenicol acetyl transferase (CAT). The CAT coding sequence was flanked by RSV-specific gene-start and gene-end motifs, the transcription signals for the viral RNA-dependent RNA polymerase. The RSV/CAT chimeric transcription cassette was inserted into the intergenic region between the G and F genes of the complete cDNA-encoded positive-sense RSV antigenome, and infectious CAT-expressing recombinant RSV was recovered. The CAT mRNA was efficiently expressed and the

levels of the G and F mRNAs were comparable to those expressed by wild type recombinant RSV. The CAT-containing and wild type viruses were similar with regard to the levels of synthesis of the major viral proteins.

5 Plasmid D46 was used for construction of cDNA encoding RSV antigenomic RNA containing the CAT gene. (Plasmids D46 and D50, the latter mentioned in Example III, are different preparations of the same antigenome cDNA.) D46 which encodes the complete, 15,223-nucleotide RSV antigenome
10 (one nucleotide longer than that of wild type RSV) and was used to produce recombinant infectious RSV described above. During its construction, the antigenome cDNA had been modified to contain four new restriction sites as markers. One of these, a *StuI* site placed in the intergenic region between the
15 G and F genes (positions 5611-5616 in the 3'-5' sequence of the wild type genome), was chosen as an insertion site for the foreign CAT gene. A copy of the CAT ORF flanked on the upstream end by the RSV GS signal and on the downstream end by the RS GE signal was derived from a previously-described RSV-
20 CAT minigenome (Collins et al., Proc. Natl. Acad. Sci. USA 88:9663-9667 (1991) and Kuo et al., J. Virol. 70: 6892-6901 (1996), incorporated by reference herein). The insertion of this RSV/CAT transcription cassette into the *StuI* site, to yield the D46/1024CAT cDNA, increased the length of the
25 encoded antigenome to a total of 15,984 nucleotides. And, whereas wild type RSV encodes ten major subgenomic mRNAs, the recombinant virus predicted from the D46/1024CAT antigenome would encode the CAT gene as an eleventh mRNA. The strategy of construction is shown in Fig. 2.

30 Producing infectious RSV from cDNA-encoded antigenomic RNA, as described above, involved coexpression in HEP-2 cells of five cDNAs separately encoding the antigenomic RNA or the N, P, L or M2(ORF1) protein, which are necessary and sufficient for viral RNA replication and transcription.
35 cDNA expression was driven by T7 RNA polymerase supplied by a vaccinia-T7 recombinant virus based on the MVA strain. The MVA-T7 recombinant virus produced infectious progeny

sufficient to cause extensive cytopathogenicity upon passage, and therefore, cytosine arabinoside, an inhibitor of vaccinia virus replication, was added 24 h following the transfection and maintained during the first six passages.

5 Two antigenome cDNAs were tested for the recovery of RSV: the D46 cDNA, and the D46/1024CAT cDNA. Each one yielded infectious recombinant RSV. Cells infected with the D46/1024CAT recombinant virus expressed abundant levels of CAT enzyme. For each virus, transfection supernatants were
10 passaged to fresh cells, and a total of eight serial passages were performed at intervals of five to six days and a multiplicity of infection of less than 0.1 PFU per cell.

The CAT sequence in the D46/1024CAT genome was flanked by RSV GS and GE signals, and thus should be expressed
15 as an additional, separate, polyadenylated mRNA. The presence of this predicted mRNA was tested by Northern blot hybridization of RNA from cells infected with D46/1024CAT virus or D46 virus at the eighth passage. Hybridization with a negative-sense CAT-specific riboprobe detected a major band
20 which was of the appropriate size to be the predicted CAT mRNA, which would contain 735 nucleotides not including poly(A). This species was completely retained by oligo(dT) latex particles, showing that it was polyadenylated. In some cases, a minor larger CAT-specific species was detected which
25 was of the appropriate size to be a G-CAT readthrough mRNA. The D46/1024CAT virus had been subjected to eight passages at low multiplicity of infection prior to the infection used for preparing the intracellular RNA. There was no evidence of shorter forms of the CAT mRNA, as might have arisen if the CAT
30 gene was subject to deletion.

Replicate blots were hybridized with negative-sense riboprobe specific to the CAT, SH, G or F gene, the latter two genes flanking the inserted CAT gene. The blots showed that
35 the expression of the subgenomic SH, G and F mRNAs was similar for the two viruses. Phosphoimager was used to compare the amount of hybridized radioactivity in each of the three RSV mRNA bands for D46/1024CAT and D46. The ratio of

radioactivity between D46/1024CAT and D46 was determined for each mRNA: SH, 0.77; G, 0.87; and F, 0.78. The deviation from unity probably indicates that slightly less RNA was loaded for D46/1024CAT versus D46, although it also is possible that the overall level of mRNA accumulation was slightly less for D46/1024CAT RSV. The demonstration that the three ratios were similar confirms that the level of expression of each of these mRNAs was approximately the same for D46/1024CAT versus D46. This, the insertion of the CAT gene between the G and F genes did not drastically affect the level of transcription of either gene.

To characterize viral protein synthesis, infected HEP-2 cells were labeled with [³⁵S]methionine, and cell lysates were analyzed by PAGE either directly or following immunoprecipitation under conditions where recovery of labeled antigen was essentially complete. Precipitation with a rabbit antiserum raised against purified RSV showed that the D46/1024CAT and D46 viruses both expressed similar amounts of the major viral proteins F₁, N, P, M, and M2. That a similar level of M2 protein was recovered for each virus was noteworthy because its gene is downstream of the inserted CAT gene. Accumulation of the F protein, which is encoded by the gene located immediately downstream of the insertion, also was examined by immunoprecipitation with a mixture of three anti-F monoclonal antibodies. A similar level of the F₁ subunit was recovered for each virus. Phosphorimager analysis of the major viral proteins mentioned above was performed for several independent experiments and showed that some sample-to-sample variability, but overall the two viruses could not be distinguished on the basis of the level of recovered proteins. Precipitation with anti-CAT antibodies recovered a single species for the D46/1024CAT but not for the D46 virus. Analysis of the total labeled protein showed that the N, P and M proteins could be detected without immunoprecipitation (although detection of the latter was complicated by its comigration with a cellular species) and confirmed that the two viruses yielded similar patterns. The position

corresponding to that of the CAT protein contained more radioactivity in the D46/1024CAT pattern compared to that of D46, as was confirmed by phosphorimager of independent experiments. This suggested that the CAT protein could be
5 detected among the total labeled proteins without precipitation, although this demonstration was complicated by the presence of a comigrating background band in the uninfected and D46-infected patterns.

RT-PCR was used to confirm the presence of the CAT
10 gene in the predicted location of the genome of recombinant RSV. Total intracellular RNA was isolated from the cell pellet of passage eight of both D46/1024CAT and D46 RSV. Two primers were chosen that flank the site of insertion, the StuI restriction endonuclease site at RSV positions 5611-5616: the
15 upstream positive-sense primer corresponded to positions 5412-5429, and the downstream negative-sense one to positions 5730-5711. The positive-sense primer was used for the RT step, and both primers were included in the PCR.

RT-PCR of the D46 virus yielded a single product
20 that corresponded to the predicted fragment of 318 nucleotides, representing the G/F gene junction without additional foreign sequence. Analysis of D46/1024CAT viral RNA yielded a single product whose electrophoretic mobility corresponded well with the predicted 1079 nucleotide fragment,
25 representing the G/F gene junction containing the inserted CAT transcription cassette. The latter PCR yielded a single major band; the absence of detectable smaller products indicated that the population of recombinant genomes did not contain a large number of molecules with a deletion in this region.
30 When PCR analysis was performed on D46/1024CAT virus RNA without the RT step, no band was seen, confirming that the analysis was specific to RNA. Thus, the RT-PCR analysis confirmed the presence of an insert of the predicted length in the predicted location in the genomic RNA of the D46/1024CAT
35 recombinant virus.

Enzyme expression was used to measure the stability of the CAT gene. Cell pellets from all of the passages

beginning with the third were tested for CAT expression. For the virus D46/1024CAT, all these assays displayed conversion of [14 C] labeled chloramphenicol into acetylated forms. To investigate stability of expression, virus from 20 or 25 individual plaques from passage three or eight, respectively, was analyzed for CAT expression. All samples were positive, and the level of expression of CAT was similar for each of the 25 isolates from passage eight, as judged by assay of equivalent aliquots of cell lysate. This demonstrated that the activity of the CAT protein encoded by each isolate remained unimpaired by mutation.

To determine plaque morphology and size, beginning with the second passage, one-eighth of the medium supernatant (i.e., 0.5 ml) harvested from each passage stage was used to infect fresh HEp-2 cells in six-well plates that were incubated under methylcellulose overlay for five to seven days. The cells were then fixed and stained by incubation with monoclonal antibodies against RSV F protein followed by a second antibody linked to horseradish peroxidase. Earlier, it had been observed that recombinant RSV produced from cDNA D46 was indistinguishable from a naturally-occurring wild type RSV isolate with regard to efficiency of plaque formation over a range of temperatures *in vitro*, and the ability to replicate and cause disease when inoculated into the respiratory tract of previously uninfected chimpanzees. Thus, the D46 recombinant RSV was considered to be a virulent wild type strain. The plaques produced by the D46 and D46/1024CAT recombinant viruses were compared by antibody staining. Plaque morphology was very similar for the two viruses, although the average diameter of the CAT-containing recombinant plaques was 90 percent of that of the D46 virus, based on measurement of thirty randomly-selected plaques for each virus.

The efficiency of replication in tissue culture of the D46 and D46/1024CAT viruses was compared in a single step growth cycle. Triplicate monolayers of cells were infected with either virus, and samples were taken at 12 h intervals

and quantitated by plaque assay. The results showed that the production of D46/1024CAT virus relative to D46 was delayed and achieved a maximum titer which was 20-fold lower.

These results show that it is possible to construct recombinant, helper-independent RSV expressing a foreign gene, in this instance the CAT gene. The recombinant RSV directed expression of the predicted polyadenylated subgenomic mRNA that encoded CAT protein, the protein being detected both by enzyme assay and by radioimmunoprecipitation. Other examples have produced RSV recombinants with the luciferase gene inserted at the same CAT site, or with the CAT or luciferase genes inserted between the SH and G genes. These viruses also exhibit reduced growth, whereas the numerous wild type recombinant viruses recovered exhibit undiminished growth. This indicates that the reduced growth indeed is associated with the inserted gene rather than being due to chance mutation elsewhere in the genome. The finding that insertion of a foreign gene into recombinant RSV reduced its level of replication and was stable during passage in vitro suggests that this provides yet another means for effecting attenuation for vaccine use. And, these results demonstrate that the methodology described herein is capable of recovering a virus that is restricted in growth.

These results also illustrate an advantage of the strategy of gene expression of the nonsegmented negative strand viruses, namely that the foreign coding sequences can be introduced as a separate transcription cassette that is expressed as a separate mRNA. The results also show that RSV can tolerate an increase of genome length of 762 nucleotides in the case of the CAT gene to a total of 15,984 nucleotides (1.05 times that of wild type RSV). The luciferase gene that was successfully recovered is almost three times longer.

The viral RNA-dependent RNA polymerases are known to have an error-prone nature due to the absence of proofreading and repair mechanisms. In RNA virus genomes, the frequency of mutation is estimated to be as high as 10^{-4} - 10^{-5} per site on average (Holland et al., Curr. Top. Microbiol. Immunol. 176:1-

20 (1992) and references therein). In the case of the recombinant D46/1024CAT RSV produced here, correct expression of the foreign gene would be irrelevant for virus replication and would be free to accumulate mutations. The passages
5 described here involved a multiplicity of infection less than 0.1 PFU per cell, and the duration of each passage level indicated that multiple rounds of infection were involved. While yields of infectious virus from RSV-infected tissue culture cells typically are low, intracellular macromolecular
10 synthesis is robust, and the poor yields of infectious virus seems to represent an inefficient step in packaging rather than low levels of RNA replication. Thus, the maintenance of CAT through eight serial passages involved many rounds of RNA replication. It was surprising that the nonessential CAT gene
15 remained intact and capable of encoding fully functional protein in each of the 25 isolates tested at the eighth passage. Also, RT-PCR analysis of RNA isolated from passage eight did not detect deletions within the CAT gene.

Because most of the antigenic difference between the
20 two RSV antigenic subgroups resides in the G glycoprotein, recombinant RSV can be constructed to express the G protein of the heterologous subgroup as an additional gene to yield a divalent vaccine. Envelope protein genes of some other
25 respiratory viruses, such as human parainfluenza 3 virus, also can be inserted for expression by recombinant RSV. Other uses include coexpression of immune modulators such as interleukin
6 to enhance the immunogenicity of infectious RSV. Other uses, such as employing modified RSV as described herein as a vector for gene therapy, are also provided.

30 Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: The Government of the United States of America, as represented by the Secretary of the Department of Health and Human Services

(B) STREET: Box OTT
(C) CITY: Bethesda
(D) STATE: Maryland
(E) COUNTRY: United States of America
(F) POSTAL CODE (ZIP): 20892
(G) TELEPHONE: (301) 496-7056
(H) TELEFAX: (301) 402-0220
(I) TELEX:

(ii) TITLE OF INVENTION: PRODUCTION OF INFECTIOUS RESPIRATORY SYNCYTIAL VIRUS FROM CLONED NUCLEOTIDE SEQUENCES

(iii) NUMBER OF SEQUENCES: 1

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(v) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: WO
(B) FILING DATE:
(C) CLASSIFICATION:

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 60/007,083
(B) FILING DATE: 27-SEP-1995

(vii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Parmelee, Steven W.
(B) REGISTRATION NUMBER: 31,990
(C) REFERENCE/DOCKET NUMBER: 15280-250-1PC

(viii) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 206-467-9600
(B) TELEFAX: 415-576-0300

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15223 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ACGSGAAAAA ATCGGTACAA CAACTTGCA TAAACAAAAA AAATGGGGCA AATAAGAATT	60
TGATAAGTAC CACTTAAATT TAACTCCCTT GGTTAGAGAT GGGCAGCAAT TCATTGAGTA	120

TGATAAAAGT	TAGATTACAA	AATTTGTTTG	ACAATGATGA	AGTAGCATTG	TTAAAAATAA	180
CATGCTATAC	TGATAAAITA	ATACATTITAA	CTAATGCTTT	GGCTAAGGCA	GTGATACATA	240
CAATCAAAT	GAATGGCATT	GTGTTTGTGC	ATGTTATTAC	AAGTAGTGAT	ATTTGCCCTA	300
ATAATAATAT	TGTAGTAAAA	TCCAAITTC	CAACAATGCC	AGTACTACAA	AATGGAGGTT	360
ATATATGGGA	AATGATGGAA	TTAACACATT	GCTCTCAACC	TAATGGTCTA	CTAGATGACA	420
ATTGTGAAAT	TAAATTCCTC	AAAAAACTAA	GTGATTCAAC	AATGACCAAT	TATATGAATC	480
AATTATCTGA	ATTACTTGGA	TTTGATCTTA	ATCCATAAAT	TATAATTAAAT	ATCAACTAGC	540
AAATCAATGT	CACTAACACC	ATTAGTTAAT	ATAAAACTTA	ACAGAAGACA	AAAAATGGGGC	600
AAATAAATCA	ATTGAGCCAA	CCCAACCATG	GACACAACCC	ACAATGATAA	TACACCACAA	660
AGACTGATGA	TCACAGACAT	GAGACCGTTG	TCACTTGAGA	CCATAATAAC	ATCACTAACC	720
AGAGACATCA	TAACACACAA	ATTATATATC	TTGATAAATC	ATGAATGCAT	AGTGAGAAAA	780
CTTGATGAAA	AGCAGGCCAC	ATTTACATTC	CTGCTCAACT	ATGAAATGAA	ACTATTACAC	840
AAAGTAGGAA	GCACATAATA	TAAAAAATAT	ACTGAATACA	ACACAAAATA	TGGCATTTC	900
CCTATGCCAA	TATTATCACT	TCATGATGGG	TTCTTAGAAT	GCATTGGCAT	TAAGCCTACA	960
AAGCATACTC	CCATAATATA	CAAGTATGAT	CTCAATCCAT	AAATTTCAAC	ACAATATTCA	1020
CACAATCTAA	AACAACAAC	CTATGCATBA	CTATACTCCA	TAGTCCAGAT	GGAGCCTGAA	1080
AATTATAGTA	ATTTAAACT	TAAGGAGAGA	TATAAGATAG	AAGATGGGGC	AAATACAACC	1140
ATGGCTCTTA	GCAAGTCAA	GTTGAATGAT	ACACTCAACA	AAGATCAACT	TCTGTCTATCC	1200
AGCAAAATACA	CCATCCAACG	GAGCACAGGA	GATAGTATTG	ATACTCCTAA	TTATGATGTG	1260
CAGAAACACA	TCAATAAGTT	ATGTGGCATG	TTATTAATCA	CAGAAGATGC	TAATCATAAA	1320
TTCACTGGGT	TAATAGGTAT	GTTATATGCG	ATGCTTAGGT	TAGGAAGAGA	AGACACCATA	1380
AAAATACTCA	GAGATGCGGG	ATATCATGTA	AAAGCAAATG	GAGTAGATGT	AACAACACAT	1440
CGTCAAGACA	TTAATGGAAA	AGAAATGAAA	TTTGAAGTGT	TAACATTGGC	AAGCTTAACA	1500
ACTGAAATTC	AAATCAACAT	TGAGATAGAA	TCTAGAAAT	CCTACAAAAA	AATGCTAAAA	1560
GAAATGGGAG	AGGTAGCTCC	AGAATACAGG	CATGACTCTC	CTGATTGTGG	GATGATAATA	1620
TTATGTATAG	CAGCATTAGT	AATAACTAAA	TTAGCAGCAG	GGGACAGATC	TGGTCTTACA	1680
GCCGTGATTA	GGAGAGCTAA	TAATGTCTTA	AAAAATGAAA	TGAAACGTTA	CAAAGGCTTA	1740
CTACCCAAGG	ACATAGCCAA	CAGCTTCTAT	GAAGTGTTTG	AAAAACATCC	CCACTTTATA	1800
GATGTTTTTG	TTCATTTTGG	TATAGCACAA	TCTTCTACCA	GAGGTGGCAG	TAGAGTTGAA	1860
GGGATTTTTG	CAGGATTGTT	TATGAATGCC	TATGGTGCAG	GGCAAGTGAT	GTTACGGTGG	1920
GGAGTCTTAG	CAAAATCAGT	TAAAAATATT	ATGTTAGSAC	ATGCTAGTGT	GCAAGCAGAA	1980
ATGGAACAAG	TTGTTGAGGT	TTATGAATAT	GCCCAAAAAAT	TGGGTGGTGA	AGCAGGATTC	2040
TACCATATAT	TGAACAACCC	AAAAGCATCA	TTATTATCTT	TGACTCAATT	TCCTCACTTC	2100
TCCAGTGTAG	TATTAGGCCA	TGCTGCTGGC	CTAGGCATAA	TGGGAGAGTA	CAGAGGTACA	2160

CCGAGGAATC	AAGATCTATA	TGATGCAGCA	AAGGCATATG	CTGAACAACT	CAAAAGAAAT	2220
GGTGATGATTA	ACTACAGTGT	ACTAGACTTG	ACAGCAGAAG	AACTAGAGGC	TATCAAAACAT	2280
CAGCTTAATC	CAAAAGATAA	TGATGTAGAG	CTTTGAGTTA	ATAAAAATG	GGGCAAAATA	2340
ATCATCATGG	AAAAGTTTGC	TCCTGAATTC	CATGGAGAAG	ATGCAACAA	CAGGGCTACT	2400
AAATTCTAG	AATCAATAAA	GGGCAAATC	ACATCACCCA	AAGATCCCAA	GA AAAAGAT	2460
AGTATCATAT	CTGTCAACTC	AATAGATATA	GAAGTAACCA	AAGAAAGCCC	TATAACATCA	2520
AATTCAACTA	TTATCAACCC	AACAATAGAG	ACAGATGATA	CTGCAGGGAA	CAAGCCCAAT	2580
TATCAAGAA	AACCTCTAGT	AAGTTTCAAA	GAAGACCTTA	CACCAAGTGA	TAATCCCTTT	2640
TCTAAACTAT	ACAAAGAAAC	CATAGAAACA	TTTGATAACA	ATGAAGAAGA	ATCCAGCTAT	2700
TCATACGAAG	AAATAAATGA	TCAGACAAAC	GATAATATAA	CAGCAAGATT	AGATAGGATT	2760
GATGAAAAAT	TAAGTGAAAT	ACTAGGAATG	CTTCACACAT	TAGTAGTGGC	AAGTGCAGGA	2820
CCTACATCTG	CTCGGGATGG	TATAAGAGAT	GCCATGGTTG	GTTTAAAGAGA	AGAAATGATA	2880
GA AAAAATCA	GAATGGAAGC	ATTAATGACC	AATGACAGAT	TAGAAGCTAT	GGCAAGACTC	2940
AGGAATGAGG	AAAGTGA AAA	GATGGCAAAA	GACACATCAG	ATGAAGTGTC	TCTCAATCCA	3000
ACATCAGAGA	AATTGAACAA	CCTATTGGAA	GGGAATGATA	GTGACAATGA	TCTATCACTT	3060
GAAGATTCTT	GATTAGTTAC	CAATCTTCAC	ATCAACACAC	AATACCAACA	GAAGACCAAC	3120
AAACTAACCA	ACCCATCAT	CCAACCAAC	ATCCATCCGC	CAATCAGCCA	AACAGCCAA	3180
AAACAACCA	GCCATCCAA	AACTAACCA	CCGAAAAAA	TCTATAATAT	AGTTACAAAA	3240
AAAGGAAGG	GTGGGGCAAA	TATGGAAACA	TACGTGAACA	AGCTTCACGA	AGGCTCCACA	3300
TACACAGCTG	CTGTTCAATA	CAATGCTTTA	GA AAAAGACG	ATGACCTGTC	ATCACTTACA	3360
ATATGGGTGC	CCATGTTCCA	ATCATCTATG	CCAGCAGATT	TACTTATAAA	AGAAGTACTG	3420
AATGTCAACA	TACTAGTGAA	ACAAATATCC	ACACCCAAAG	GACCTTCACT	AAGAGTCATG	3480
ATAAACTCAA	GAAGTGCAGT	GCTAGCACAA	ATGCCACGAA	AATTTACCAT	ATGCGCTAAT	3540
GTGTCCCTGG	ATGAAGAAG	CAAACTAGCA	TATGATGTAA	CCACACCCCTG	TGAATCAAG	3600
GCATGTAGTC	TAACATGCCCT	AAAATGAAA	AAATGTTTGA	CTACAGTTAA	AGATCTCACT	3660
ATGAAGACAC	TCAACCTAT	ACATGATATT	ATTGCTTTAT	GTGAATTTGA	AAACATAGTA	3720
ACATCAAAA	AAGTCATAAT	ACCAACATAC	CTAGATCCA	TCAGTGTGAG	AAATAAGAT	3780
CTGAACACAC	TTGAAAATAT	AACAACCACT	GAATTCAAAA	ATGCTATCAC	AAATGC AAAA	3840
ATCATCCCTT	ACTCAGGATT	ACTATTAGTC	ATCACAGTGA	CTGACAACAA	AGGAGCATTC	3900
AAATACATAA	AGCCACAAAG	TCAATTCTATA	GTAGATCTTG	GAGCTTACCT	AGAAAAGGAA	3960
AGTATATATT	ATGTTACCAC	AAATTGGAAG	CACACAGCTA	CAGGATTGTC	AATCAAAACC	4020
ATGGAAGATT	AACCTTTTTT	CTCTACATCA	GTGTGTTAAT	TCATACAAAC	TTTCTACCTA	4080
CATTCTTCAC	TTCAACATCA	CAATCAAAA	CACCTCTGTG	TTCAACCAAT	CAACAAAAC	4140
TTATCTGAAG	TCCACAGTCA	TCCCAAGTCA	TTGTTTATCA	GATCTAGTAC	TCAATAAGT	4200

TAATAAAAA	TATACACATG	GGGCAATAA	TCATTGGAGG	AAATCCAAC	AATCACAATA	4260
TCTGTTAACA	TAGACAAGTC	CACACACCAT	ACAGAATCAA	CCAAATGGAAA	ATACATCCAT	4320
AACAATAGAA	TTCTCAAGCA	AATTCTGGCC	TTACTTTACA	CTAATACACA	TGATCACAAC	4380
AATAATCTCT	TTGCTAATCA	TAATCTCCAT	CATGATTGCA	ATACTAAACA	AACCTTGTGA	4440
ATATAACGTA	TTCCATAACA	AAACCTTTGA	GTACCAAGA	GCTCGAGTCA	ACACATAGCA	4500
TTCTCAATC	CAACAGCCCA	AAACAGTAAC	CTTGCAITTA	AAAATGAACA	ACCCTACCT	4560
CTTTACAACA	CCTCATTAA	ATCCACCAT	GCAACCACCT	ATCCATACTA	TAAAGTAGTT	4620
AATTAATAAT	AGTCATAACA	ATGAACTAGG	ATATCAAGAC	TAACAATAAC	ATTGGGGCAA	4680
ATGCAAAACAT	GTCCAAAAAC	AAGGACCAAC	GCACCGCTAA	GACATTAGAA	AGGACCTGGG	4740
ACACTCTCAA	TCATTATTATA	TTCATATCAT	CGTGCTTATA	TAAGTTAAAT	CTTAAATCTG	4800
TAGCACAAAT	CACATTATCC	ATTCTGGCAA	TGATAATCTC	AACTTCACTT	ATAATTGCGAG	4860
CCATCATATT	CATAGCCTCG	GCAAACCACA	AAGTCACACC	AACAACCTGCA	ATCATACAAG	4920
ATGCAACAAG	CCAGATCAAG	AACACAACCC	CAACATACCT	CACCAGAAT	CCTCAGCTTG	4980
GAATCAGTCC	CTCTAATCCG	TCTGAAATTA	CATCACAAT	CACCACCATA	CTAGCTTCAA	5040
CAACACCAGG	AGTCAAGTCA	ACCCTGCAAT	CCACACAGT	CAAGACCAAA	AACACAACAA	5100
CAACTCAAAC	ACAACCCAGC	AAGCCACCA	CAAAACACG	CCAAAACAA	CCOCCAAGCA	5160
AACCCAATAA	TGATTTTTCAC	TTTGAAGTGT	TCAACTTTGT	ACCCTGCAGC	ATATGCAGCA	5220
ACAAATCAAC	CTGCTGGGCT	ATCTGCAAAA	GAATACCAAA	CAAAAAACCA	GGAAAGAAAA	5280
CCACTACCAA	GCCACAAAA	AAACCAACCC	TCAAGACAA	CAAAAAAGAT	CCCAACCTC	5340
AAACCACATA	ATCAAAGGAA	GTACCCACCA	CCAAGCCAC	AGAAGAGCCA	ACCATCAACA	5400
CCACCAAAAC	AAACATCATA	ACTACACTAC	TCACCTCCAA	CACCACAGGA	AATCCGAAC	5460
TCACAAGTCA	AATGGAAACC	TTCCACTCAA	CTTCTCCGA	AGGCAATCCA	AGCCCTTCTC	5520
AAGTCTCTAC	AACATCCGAG	TACCATCAC	AACCTTCATC	TCCACCCAA	ACACCAGCC	5580
AGTAGTTACT	TAAAAACATA	TTATCACAAA	AGGCCTTGAC	CAACTTAAAC	AGAATCAAAA	5640
TAAACTCTGG	GGCAATAAAC	AATGGAGTTG	CTAATCCTCA	AAGCAATGC	AATTACCACA	5700
ATCCTCACTG	CAGTCACATT	TTGTTTTGCT	TCTGGTCAAA	ACATCACTGA	AGAATTTTAT	5760
CAATCAACAT	GCAGTGCAGT	TAGCAAGGC	TATCTTAGTG	CTCTGAGAAC	TGGTTGGTAT	5820
ACCAGTGTTA	TAACATAGAA	ATTAAATTA	ATCAAGAAA	ATAAGTGTA	TGGAACAGAT	5880
GCTAAGGTAA	AATTGATAAA	ACAAGAATTA	GATAATATA	AAAATGCTGT	AACGAAGATTG	5940
CAGTTGCTCA	TGCAAGGAC	ACAAGCAACA	AACATCGAG	CCAGAAGAGA	ACTACCAAGG	6000
TTTATGAATT	ATACACTCAA	CAATGCCAAA	AAAACCAAT	TACATTTAAG	CAAGAAAAGG	6060
AAAAGAAGAT	TTCTTGTTTT	TTTGTAGGT	GTGGATCTG	CAATCGCCAG	TGGCGTTGCT	6120
GTATCTAAGG	TCCTGCACCT	AGAAGGGGAA	GTGAACAAGA	TCAAAGTGC	TCTACTATCC	6180
ACAAACAAGG	CTGTAGTCAG	CTTATCAAAT	GGAGTTAGTG	TTTTAACCAG	CAAAGTGTTA	6240

GACCTCAAAA	ACTATATAGA	TAAACAATTG	TTACCTATTG	TGAACAAGCA	AAGCTGCAGC	6300
ATATCAAAATA	TAGAAACTGT	GATAGAGTTC	CAACAAAAGA	ACAACAGACT	ACTAGAGATT	6360
ACCAGGGAAAT	TTAGTGTATA	TGCAGGCGTA	ACTACACCTG	TAAGCACTTA	CATGTAACT	6420
AATAGTGAAT	TATTGTCATT	AATCAATGAT	ATGCCTATAA	CAAATGATCA	GAATAAGTTA	6480
ATGTCCAACA	ATGTTCAAAT	AGTTAGACAG	CAAAGTTACT	CTATCATGTC	CATAATAAAA	6540
GAGGAAGTCT	TAGCATATGT	AGTACAATTA	CCACTATATG	GTGTTATAGA	TACACCTGT	6600
TGGAAACTAC	ACACATCCCC	TCTATGTACA	ACCAACACAA	AAGAAGGGTC	CAACACTCTG	6660
TTAACAAGAA	CTGACAGAGG	ATGGTACTGT	GACAAATGCAG	GATCAGTATC	TTTCTTCCCA	6720
CAAGCTGAAA	CATGTAAAGT	TCAATCAAAT	CGAGTATTTT	GTGACACAAT	GAACAGTTTA	6780
ACATTACCAA	GTGAAGTAAA	TCTCTGCAAT	GTTGACATAT	TCAACCCCAA	ATATGATTGT	6840
AAAATTATGA	CTTCAAAAAC	AGATGTAAGC	AGCTCCGTTA	TCACATCTCT	AGGAGCCATT	6900
GTGTCATGCT	ATGGCAAAAC	TAAATGTACA	GCATCCRAAT	AAAATCGTGG	AATCATAAAG	6960
ACATTTTCTA	ACGGGTGCGA	TTATGTATCA	AATAAAGGGG	TGGACACTGT	GTCTGTAGST	7020
AACACATTAT	ATTATGTAAA	TAAGCAAGAA	GGTAAAAGTC	TCTATGTAAA	AGGTGAACCA	7080
ATAATAAATT	TCTATGACCC	ATTAGTATTC	CCCTCTGATG	AATTTGATGC	ATCAATATCT	7140
CAAGTCAACG	AGAAGATTAA	CCAGAGCCTA	GCAATTTATC	GTAAATCCGA	TGAATTATTA	7200
CATAATGTAA	ATGCTGTATA	ATCCACCACA	AATATCATGA	TAACTACTAT	AATTATAGTG	7260
ATTATAGTAA	TATTGTTATC	ATTAAATGCT	GTTGGACTGC	TCTTATACGT	TAAGGCCAGA	7320
AGCACACCAG	TCACACTAAG	CAAAGATCAA	CTGAGTGGTA	TAAATAATAT	TGCATTTAGT	7380
AACATAATAA	AAATAGCACC	TAATCATGTT	CTTACAATGG	TTTACTATCT	GCTCATAGAC	7440
AACCCATCTG	TCATTGGATT	TTCTTAAAAAT	CTGAACCTCA	TCGAAACTCT	CATCTATAAA	7500
CCATCTCACT	TACACTATTT	AAGTAGATTG	CTAGTTTATA	GTTATATAAA	ACACAATTGC	7560
ATGCCAGATT	AACCTTACCAT	CTGTAAAAAT	GAAACTGGG	GCAATATGT	CACGAAGGAA	7620
TCCTTGCAAA	TTTGAAATTC	GAGGTCAATT	CTTAAATGGT	AAGAGGTGTC	ATTTTATGCA	7680
TAATTATTTT	GAATGGCCAC	CCCATGCACT	GCTTGTAAGA	CAAACTTTTA	TGTTAAACAG	7740
AATACTTAAG	TCTATGGATA	AAAGTATAGA	TACCTTATCA	GAATAAGTG	GAGCTGCAGA	7800
GTTGACAGAA	ACGAAGAGT	ATGCTCTTGG	TGTAGTTGGA	GTGCTAGAGA	GTTATATAGG	7860
ATCAATAAAC	AATATAACTA	AACAATCAGC	ATGTGTTGCC	ATGAGCAAAC	TCCTCACTGA	7920
ACTCAATAGT	GATGATATCA	AAAAGCTGAG	GGACAATGAA	GAGCTAAATT	CACCCAAGAT	7980
AAGAGTGTAC	AATACTGTCA	TATCATATAT	TGAAGCAAC	AGGAAAAACA	ATAAACAAAC	8040
TATCCATCTG	TTAAAAAGAT	TGCCAGCAGA	CGTATTGAAG	AAAACCATCA	AAACACATT	8100
GGATATCCAT	AAGAGCATAA	CCATCAACAA	CCCAAAAGAA	TCAACTGTTA	GTGATACAAA	8160
TGACCATGCC	AAAAATAATG	ATACTACCTG	ACAAATATCC	TTGTAGTATA	ACTTCCATAC	8220
TAATAACAAG	TAGATGTAGA	GTTACTATGT	ATAATCAAAA	GAACACACTA	TATTTCAATC	8280

AAAACAACCC	AAATAACCAT	ATGTACTCAC	CGAATCAAAC	ATTCAATGAA	ATCCATTGGA	8340
CCTCTCAAGA	ATTGATTGAC	ACAATTCAAA	ATTTTCTACA	ACATCTAGGT	ATTATTGAGG	8400
ATATATATAC	AATATATATA	TTAGTGTCA	AACACTCAAT	TCTAACACTC	ACCACATCGT	8460
TACATTATTA	ATTCAAACAA	TTCAAGTTGT	GGGACAAAAT	GGATCCCATT	ATTAATGGAA	8520
ATTCTGCTAA	TGTTTATCTA	ACCGATAGTT	ATTTAAAAGG	TGTTATCTCT	TTCTCAGAGT	8580
GTAATGCTTT	AGGAAGTTAC	ATATTCCAAT	GTCTTATCT	CAAAAATGAT	TATACCAACT	8640
TAATTAGTAG	ACAAAATCCA	TTAATAGAAC	ACATGAATCT	AAAGAAACTA	AATATAACAC	8700
AGTCCTTAAT	ATCTAAGTAT	CATAAAGGTG	AAATAAAATT	AGAAGAACCT	ACTTATTTTC	8760
AGTCATTACT	TATGACATAC	AAGAGTATGA	CCTCGTCAGA	ACAGATTGCT	ACCACTAATT	8820
TACTTAAAAA	GATAATAAGA	AGAGCTATAG	AAATAAGTGA	TGTCAAAGTC	TATGCTATAT	8880
TGAATAAACT	AGGGCTTAAA	GAAAAGGACA	AGATTAAATC	CAACAATGGA	CAAGATGAAG	8940
ACAACCTCAGT	TATTACGACC	ATAATCAAAG	ATGATATACT	TTCAGCTGTT	AAAGATAATC	9000
AATCTCATCT	TAAAGCAGAC	AAAAATCACT	CTACAAAACA	AAAAGACACA	ATCAAAAACA	9060
CACCTCTTGAA	GAAATTGATG	TGTTCAATGC	AACATCCTCC	ATCATGGTTA	ATACATTGGT	9120
TTAACCTTATA	CACAAAATTA	AACAACATAT	TAACACAGTA	TCGATCAAAT	GAGGTAAAAA	9180
ACCATGGGTT	TACATTGATA	GATAATCAAA	CTCTTAGTGG	ATTTCAAATT	ATTTTGAACC	9240
AATATGGTTG	TATAGTTTAT	CATAAGGAAC	TCAAAGAAT	TACTGTGACA	ACCTATAATC	9300
AATCTTGTAC	ATGGAAGAGT	ATTAGCCTTA	GTAGATTAAG	TGTTTGTTTA	ATTACATGGA	9360
TTAGTAACCTG	CTTGAACACA	TTAAATAAAA	GCCTTAGGCT	AAGATGCGGA	TTCAATAATG	9420
TTATCTTGAC	ACAACTATT	CTTTATGGAG	ATTGTATACT	AAAGCTATT	CACAATGAGG	9480
GGTTCTACAT	AATAAAGAG	GTAGAGGGAT	TTATTATGTC	TCTAATTTTA	AATATAACAG	9540
AAGAAGATCA	ATTCAGAAAA	CGATTTTATA	ATAGTATGCT	CAACAACATC	ACAGATGCTG	9600
CTAATAAAGC	TCAGAAAAAT	CTGCTATCAA	GAGTATGTCA	TACATTATTA	GATAAGACAG	9660
TGTCOGATAA	TATAATAAAT	GGCAGATGGA	TAATTCTATT	AAGTAAGTTC	CTTAAATTAA	9720
TTAAGCTTGC	AGGTGACAA	AACCTTAACA	ATCTGAGTGA	ACTATATTTT	TTGTTACAGAA	9780
TATTTGGACA	CCCAATGGTA	GATGAAAGAC	AAGCCATGGA	TGCTGTTAAA	ATTAATTGCA	9840
ATGAGACCAA	ATTTTACTTG	TTAAGCAGTC	TGAGTATGTT	AAGAGGTGCC	TTTATATATA	9900
GAATTATAAA	AGGGTTTGTA	AATAATTACA	ACAGATGGCC	TACTTTAAGA	AATGCTATTG	9960
TTTTACCCTT	AAGATGGTTA	ACTTACTATA	AACATAACAC	TTATCCTTCT	TTGTTGGAAC	10020
TTACAGAAAG	AGATTTGATT	GTGTTATCAG	GACTACGTTT	CTATCGTGAG	TTTCGGTTGC	10080
CTAAAAAAGT	GGATCTTGAA	ATGATTATAA	ATGATAAAGC	TATATCACCT	CCTAAAAATT	10140
TGATATGGAC	TAGTTTCCTC	AGAAATTACA	TGCCATCACA	CATACAAAAC	TATATAGAAC	10200
ATGAAAAAAT	AAAATTTTCC	GAGAGTGATA	AATCAAGAAG	AGTATTAGAG	TATTATTTTAA	10260
GAGATAACAA	ATTCAATGAA	TGTGATTTAT	ACAACGTGTG	AGTTAATCAA	AGTTATCTCA	10320

ACAACCCCTAA	TCATGTGGTA	TCATTGACAG	GCAAAGAAAG	AGAACTCAGT	GTAGGTAGAA	10380
TGTTTGC AAT	GCAACCGGGA	ATGTTGACAG	AGGTTCAAAT	ATTGGCAGAG	AAAATGATAG	10440
CTGAAAACAT	TTTACAATTC	TTTCCTGAAA	GTCTTACAAG	ATATGGTGTAG	CTAGAACTAC	10500
AAAAAATATT	AGAAGTGA	GCAGGAATAA	GTAACAATTC	AAATCGTAC	AATGATAATT	10560
ACAACAATTA	CATTAGTAAG	TGCTCTATCA	TCACAGATCT	CAGCAAATTC	AATCAAGCAT	10620
TTGATATGA	AACGTCATGT	ATTTGTAGTG	ATGTGCTGGA	TGAACTGCAT	GGTGTACAAT	10680
CTCTATTTTC	CTGGTTACAT	TTAACTATTC	CTCATGTCC	AATAATATGC	ACATATAGGC	10740
ATGCAACCCC	CTATATAGGA	GATCATATTG	TAGATCTTAA	CAATGTAGAT	GAACAAAGTG	10800
GATTATATAG	ATATCACATG	GGTGGCATCG	AAGGTTGGTG	TCAAAAATA	TGGACCATAG	10860
AAGCTATATC	ACTATTGGAT	CTAATATCTC	TCAAAGGGAA	AATTCTCAAT	ACTGCTTTAA	10920
TTAATGGTGA	CAATCAATCA	ATAGATATAA	GCAAACCAAT	CAGACTCATG	GAAGGTCAAA	10980
CTCATGCTCA	AGCAGATTAT	TTGCTAGCAT	TAAATAGCCT	TAAATTACTG	TATAAAGAGT	11040
ATGCAGGCAT	AGGCCACAAA	TTAAAAGGAA	CTGAGACTTA	TATATCAAGA	GATATGCAAT	11100
TTATGAGTAA	AACAATTCAA	CATAACGGTG	TATATTACCC	AGCTAGTATA	AAGAAAGTCC	11160
TAAGAGTGGG	ACCGTGGATA	AACACTATAC	TTGATGATTT	CAAGTGAGT	CTAGAATCTA	11220
TAGGTAGTTT	GACACAAGAA	TTAGAATATA	GAGGTGAAAG	TCTATTATGC	AGTTTAAATAT	11280
TTAGAAATGT	ATGGTTATAT	AATCAGATTG	CTCTACAATT	AAAAATCAT	GCATTATGTA	11340
ACAATAAACT	ATATTTTGGAC	ATATTAAAGG	TTCTGAAACA	CTTAAAAACC	TTTTTTAATC	11400
TTGATATAT	TGATACAGCA	TTAACATTGT	ATATGAATTT	ACCCATGTTA	TTTGGTGGTG	11460
GTGATCCCAA	CTTGTTATAT	CGAAGTTTCT	ATAGAAGAAC	TCCTGACTTC	CTCACAGAGG	11520
CTATAGTTCA	CTCTGTGTTT	ATACCTAGTT	ATTATACAAA	CCATGACTTA	AAAGATAAAC	11580
TTCAAGATCT	GTCAGATGAT	AGATTGAATA	AGTTCTTAAC	ATGCATAATC	ACGTTTGACA	11640
AAAACCCCTAA	TGCTGAATTC	GTAACATTGA	TGAGAGATCC	TCAAGCTTTA	GGGTCTGAGA	11700
GACAAGCTAA	AATTACTAGC	GAAATCAATA	GACTGGCAGT	TACAGAGGTT	TTGAGTACAG	11760
CTCCAAACAA	AATATTCTCC	AAAAGTGCAC	AACATTATAC	TACTACAGAG	ATAGATCTAA	11820
ATGATATTAT	GCAAAATATA	GAACCTACAT	ATCCTCATGG	GCTAAGAGTT	GTTTATGAAA	11880
GTTTACCCTT	TTATAAAGCA	GAGAAAATAG	TAAATCTTAT	ATCAGGTACA	AAATCTATAA	11940
CTAACATACT	GGAAAAAACT	TCTGCCATAG	ACTTAACAGA	TATTGATAGA	GCCACTGAGA	12000
TGATGAGGAA	AAACATAACT	TTGCTTATAA	GGATACTTCC	ATTGGATTGT	AACAGAGATA	12060
AAAGAGAGAT	ATTGAGTATG	GAAAACCTAA	GTATTACTGA	ATTAAGCAAA	TATGTTAGGG	12120
AAAGATCTTG	GTCTTTATCC	AATATAGTTG	GTGTTACATC	ACCCAGTATC	ATGTATACAA	12180
TGGACATCAA	ATATACTACA	AGCACTATAT	CTAGTGGCAT	AATTATAGAG	AAATATAATG	12240
TTAACAGTTT	AACACGTGGT	GAGAGAGGAC	CCACTAAACC	ATGGGTTGGT	TCATCTACAC	12300
AAGAGAAAAA	AACAATGCCA	GTTTATAATA	GACAAGTCTT	AACCAAAAAA	CAGAGAGATC	12360

AAATAGATCT	ATTAGCAAAA	TIGGATTGGG	TGTATGCATC	TATAGATAAC	AAGGATGAAT	12420
TCATGGAAGA	ACTCAGCATA	GGAAACCTTG	GGTTAACATA	TGAAAAGGCC	ANGAAATTAT	12480
TTCCACAATA	TTTAAGTGTG	AATTATTTCG	ATCGCCTTAC	AGTCAGTAGT	AGACCATTGT	12540
AATTCCTCGC	ATCAATACCA	GCTTATAGAA	CAACAAATTA	TCACCTTGAC	ACTAGCCCTA	12600
TTAATCGCAT	ATTACAGAA	AAGTATGGTG	ATGAAGATAT	TGACATAGTA	TTCCAAAACT	12660
GTATAAGCTT	TGGCCTTAGT	TTAATGTCAG	TAGTAGAACA	ATTACTAAT	GTATGTCCTA	12720
ACAGAATTAT	TCTCATACCT	AAGCTTAATG	AGATACATTT	GATGAAACCT	CCCATTATCA	12780
CAGGTGATGT	TGATATTCCAC	AAGTTAAAAAC	AAGTGATACA	AAAACAGCAT	ATGTTTTTAC	12840
CAGACAAAT	AAGTTTGACT	CAATATGTGG	AATTATTCTT	AAGTAATAAA	ACACTCAAAT	12900
CTGGATCTCA	TGTTAATTCT	AATTTAATAT	TGGCACATAA	AATATCTGAC	TATTTTCATA	12960
ATACTTACAT	TTTAAGTACT	AATTTAGCTG	GACATTGGAT	TCTGATTATA	CAACTTATGA	13020
AAGATTCTAA	AGGTATTTTT	GAAAAAGATT	GGGAGAGGGG	ATATATAACT	GATCATATGT	13080
TATTAAATT	GAAAGTTTTT	TTCATGCTT	ATAAGACCTA	TCTCTTGTT	TTTCATAAAG	13140
GTTATGGCAA	AGCAAGCTG	GAGTGTGATA	TGAACACTTC	AGATCTTCTA	TGTGTATTGG	13200
AATTAATAGA	CAGTAGTTAT	TGGAAGTCTA	TGTCTAAGGT	ATTTTITAGAA	CAAAAAGTTA	13260
TCAAATACAT	TCTTAGCCAA	GATGCAAGTT	TACATAGAGT	AAAAGGATGT	CATAGCTTCA	13320
AATTATGGTT	TCTTAAACGT	CTTAATGTAG	CAGAATTCAC	AGTTTGCCCT	TGGSTTGTTA	13380
ACATAGATTA	TCATCCAACA	CATATGAAAG	CAATATTAAAC	TTATATAGAT	CTTGTTAGAA	13440
TGGGATTGAT	AAATATAGAT	AGAATACACA	TTAAAAATRA	ACACAATTC	AATGATGAAT	13500
TTTATACTTC	TAATCTCTTC	TACATTAATT	ATAACTTCTC	AGATAATACT	CATCTATTAA	13560
CTAAACATAT	AAGGATTGCT	AATCTGGAAT	TAGAAAATAA	TTACAACAAA	TTATATCATC	13620
CTACACCAGA	AACCTTAGAG	AATATACTAG	CCAATCCGAT	TAAAAGTAAT	GACAAAAAGA	13680
CACCTGAATGA	CTATTGTATA	GGTAAAAATG	TTGACTCAAT	AATGTTACCA	TGTGTTATCTA	13740
ATAAGAAGCT	TATTAATCG	TCTGCAATGA	TTAGAACCAA	TTACAGCAAA	CAAGATTGTG	13800
ATAATTTATT	CCCTATGSET	GTGATTGATA	GAATTATAGA	TCATTCAGGC	AATACAGCCA	13860
AATCCAACCA	ACTTTACACT	ACTACTTCCC	ACCAAATATC	CTTAGTGCAC	AATAGCACAT	13920
CACTTTACTG	CATGCTTCTT	TGGCATCATA	TTAATAGATT	CAATTTTGTA	TTTAGTTCTA	13980
CAGGTGTGTA	AATTAGTATA	GAGTATATTT	TAAAAGATCT	TAAAATTAAA	GATCCCAATT	14040
GTATAGCATT	CATAGGTGAA	GGAGCAGGGA	ATTTATTATT	GCGTACAGTA	GTGGAACCTC	14100
ATCCTGACAT	AAGATATATT	TACAGAAGTC	TGAAAGATTG	CAATGATCAT	AGTTTACCTA	14160
TTGAGTTTTF	AAGGCTGTAC	AATGGACATA	TCAACATTGA	TTATGGTGAA	AAITTGACCA	14220
TTCTGCTAC	AGATGCAACC	AACAACATTC	ATTGGTCTTA	TTTACATATA	AAGTTTGTCTG	14280
AACCTATCAG	TCITTTTGTG	TGTGATGCCG	AATTGTCTGT	AACAGTCAAC	TGGAGTAAAA	14340
TTATAATAGA	ATGGAGCAAG	CATGTAAGAA	AGTGCAAGTA	CTGTTCTCTA	GTTAATAAAT	14400

GTATGTTAAT	AGTAAATAT	CATGCTCAAG	ATGATATTGA	TTTCAAATTA	GACAATATAA	14460
CTATATTAAA	AAC TTATGTA	TGCTTAGGCA	GTAAGTTAAA	GGGATCGGAG	GTTTACTTAG	14520
TCCTTACAAT	AGGTCTCGG	AATATATTCC	CAGTATTTAA	TG TAGTACAA	AATGCTAAAT	14580
TGATACTATC	AAGAACCRAA	AATTT CATCA	TGCCTAAGAA	AGCTGATAAA	GAGTCTATTG	14640
ATGCAAAATAT	TAAAAGTTTG	ATACCCCTTC	TTTGTTACCC	TATAACRAAA	AAAGGAATTA	14700
ATACTGCATT	GTCAAACTA	AAGAGTGTG	TTAGTGGAGA	TATACTATCA	TATTC TATAG	14760
CTGGACGTAA	TGAAGTTTTC	AGCAATAAAC	TTATAAATCA	TAAGCATATG	AACATCTTAA	14820
AATGGTTCAA	TCATGTTTTA	AATTT CAGAT	CAACAGAACT	AACTATAAC	CATTTATATA	14880
TGGTAGAATC	TACATATCCT	TACCTAAGTG	AATTGTTAAA	CAGCTTGACA	ACCAATGAAC	14940
TTAAAAAACT	GATTAAAAATC	ACAGGTAGTC	TGTTATACAA	CTTTCATAAT	GAATAATGAA	15000
TAAAGATCTT	ATAATAAAAA	TTCCCATAGC	TATACACTAA	CACTGTATTC	AATTATAGTT	15060
ATTAAAAAAT	AAAAATCATA	TAATTTTTTA	AATAACITTT	AGTGAAC TAA	TCCTAAAGTT	15120
ATCATTTTAA	TCITGSGAGGA	ATAAAATTTAA	ACCCTAATCT	AATTGGTTTA	TATGTGTATT	15180
AACTAAATTA	CGAGATATTA	GTTTTTGACA	CTTTTTTCT	CGT		15223

WHAT IS CLAIMED IS:

1. An isolated infectious RSV particle which comprises a recombinant RSV genome or antigenome, a
5 nucleocapsid (N) protein, a nucleocapsid phosphoprotein (P), a large (L) polymerase protein, and an RNA polymerase elongation factor.
2. The isolated infectious RSV particle of claim 1,
10 wherein the RNA polymerase elongation factor is M2 (ORF1) protein of RSV.
3. The isolated infectious RSV particle of claim 1, which is a subviral particle.
- 15 4. The isolated infectious RSV particle of claim 1, which is a virus.
5. The isolated infectious RSV particle of claim 3,
20 wherein the particle comprises an RSV antigenome.
6. The isolated infectious RSV particle of claim 1, wherein the particle comprises a recombinant RSV antigenome.
- 25 7. The isolated infectious RSV particle of claim 3, wherein the particle comprises an RSV genome.
8. The isolated infectious RSV particle of claim 1, wherein the particle comprises a recombinant RSV genome.
- 30 9. The isolated infectious RSV virus of claim 1, which is a human RSV.
10. The isolated infectious RSV virus of claim 1,
35 which is a bovine or murine RSV.

11. The isolated infectious RSV of claim 1, having a genome or antigenome that is a chimera of two or more different RSV genomes.

5 12. The isolated infectious RSV virus of claim 11, wherein the chimeric genome or antigenome comprises nucleotide sequences from human and bovine RSV.

10 13. A method for producing an infectious RSV particle from one or more isolated polynucleotide molecules encoding said RSV, comprising:
 coexpressing in a cell or cell-free lysate an expression vector which comprises an isolated polynucleotide molecule encoding a RSV genome or antigenome and an expression
15 vector which comprises one or more isolated polynucleotide molecules that encodes N, P, L and RNA polymerase elongation factor proteins, thereby producing an infectious RSV particle.

20 14. The method of claim 13, wherein the RSV genome or antigenome and the N, P, L and RNA polymerase elongation factor proteins are expressed by the same expression vector.

25 15. The method of claim 13, wherein the expression vector encoding the RSV genome or antigenome and the expression vector encoding the N, P, L and RNA polymerase elongation factor proteins are different.

30 16. The method of claim 13, wherein the N, P, L and RNA polymerase elongation factor proteins are encoded on two or more different expression vectors.

 17. The method of claim 16, wherein the N, P, L and RNA polymerase elongation factor proteins are each encoded on different expression vectors.

18. The method of claim 13, wherein the isolated polynucleotide molecule that encodes a RSV genome or antigenome is cDNA.

5 19. The method of claim 13, wherein the infectious RSV particle is a virus.

20. The method of claim 19, wherein the polynucleotide molecule encoding a RSV genome or antigenome is
10 from a human, bovine or murine RSV sequence.

21. The method of claim 20, wherein the polynucleotide molecule encoding a RSV genome or antigenome is a chimera of a human RSV strain sequence and at least one non-
15 human RSV sequence.

22. The method of claim 13, wherein the polynucleotide molecule encoding the RSV genome or antigenome encodes the sequence of a wild-type RSV strain.
20

23. The method of claim 13, wherein the polynucleotide molecule encodes an RSV genome or antigenome that has been modified from a wild-type RSV strain by a nucleotide insertion, rearrangement, deletion or substitution.
25

24. The method of claim 23, wherein the modification encodes a phenotypic alteration.

25. The method of claim 24, wherein the polynucleotide molecule encodes a genome or antigenome of a nonhuman RSV.
30

26. The method of claim 24, wherein the phenotypic alteration results in attenuation, temperature-sensitivity, cold-adaptation, small plaque size or host range restriction.
35

27. The method of claim 23, wherein the polynucleotide encodes a genome or antigenome of a nonhuman RSV virus or is a chimera of a nonhuman RSV and at least one other RSV or human or nonhuman origin.

5

28. The method of claim 13, wherein the polynucleotide molecule encodes an RSV genome or antigenome of a RSV human vaccine strain that has been modified by a nucleotide insertion, deletion or substitution.

10

29. The method of claim 28, wherein the modification encodes a phenotypic alteration.

30. The method of claim 29, wherein the phenotypic alteration results in attenuation, temperature-sensitivity, cold-adaptation, small plaque size or host range restriction.

15

31. The method of claim 29, wherein the phenotypic alteration is a change in an immunogenic epitope of RSV.

20

32. The method of claim 23, wherein the polynucleotide molecule that encodes an RSV genome or antigenome of an RSV strain has been modified by inserting a nucleotide sequence that encodes a cytokine or a T-helper epitope.

25

33. The method of claim 23, wherein the polynucleotide molecule that encodes an RSV genome or antigenome of an RSV strain has been modified by inserting a nucleotide sequence encoding a restriction site marker.

30

34. The method of claim 23, wherein the polynucleotide molecule that encodes an RSV genome or antigenome of an RSV strain has been modified by inserting a nucleotide sequence encoding a G protein of an RSV subgroup different from that of said RSV strain.

35

35. The method of claim 23, wherein the polynucleotide molecule that encodes an RSV genome or antigenome of an RSV strain has been modified by inserting a nucleotide sequence encoding a protein of a microbial pathogen capable of eliciting a protective immune response.

36. The method of claim 13, wherein at least one of the viral proteins is supplied by coinfection with RSV.

37. A cell or cell-free lysate containing an expression vector which comprises an isolated polynucleotide molecule encoding a RSV genome or antigenome and an expression vector which comprises one or more isolated polynucleotide molecules that encodes N, P, L and RNA polymerase elongation factor proteins of RSV, whereby upon expression said RSV genome or antigenome and N, P, L, and RNA polymerase elongation factor proteins combine to produce an infectious RSV particle.

38. The cell or lysate of claim 37, wherein the RSV genome or antigenome and the N, P, L and RNA polymerase elongation factor proteins are encoded by the same expression vector.

39. The cell or lysate of claim 37, wherein the expression vector encoding the RSV genome or antigenome and the expression vector encoding the N, P, L and RNA polymerase elongation factor proteins are different.

40. The cell or lysate of claim 37, wherein the N, P, L and RNA polymerase elongation factor proteins are encoded on two or more expression vectors.

41. The cell or lysate of claim 40, wherein the N, P, L and RNA polymerase elongation factor proteins are each encoded on different expression vectors.

42. The cell or lysate of claim 37, wherein the infectious RSV particle is a virus.

43. The cell or lysate of claim 37, wherein the
5 polynucleotide molecule encoding a RSV genome or antigenome is a human, bovine or murine RSV sequence.

44. An isolated polynucleotide molecule which
comprises an operably linked transcriptional promoter, a
10 polynucleotide sequence encoding an RSV genome or antigenome,
and a transcriptional terminator.

45. The isolated polynucleotide molecule of claim
44, wherein the polynucleotide sequence encoding an RSV genome
15 or antigenome is a human RSV sequence.

46. The isolated polynucleotide molecule of claim
45, wherein the polynucleotide sequence encoding a human RSV
genome or antigenome is SEQ ID NO:1.

20 47. The isolated polynucleotide molecule of claim
44, wherein the polynucleotide encodes a genome or antigenome
of a nonhuman RSV virus or is a chimera of a nonhuman RSV and
at least one other RSV of human or nonhuman origin.

25

Fig. 1A





Fig. 1B



INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/15524

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N7/04 C07K14/135

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 92 07940 A (SAMAL, SIBA, KUMAR) 14 May 1992 see the whole document	1-3, 13, 17, 41, 44-47
A	---	
Y	WO 94 08022 A (COMMONWEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANIZATION) 14 April 1994 see abstract see page 7, line 9 - page 8, line 19 see page 10, line 10 - page 12, line 16 ---	1-3, 13, 17, 41, 44-47

	--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family numbers are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "Z" document member of the same patent family

Date of the actual completion of the international search

28 January 1997

Date of mailing of the international search report

26.02.97

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel.: (+31-70) 340-3040, Tlx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Panzica, G

INTERNATIONAL SEARCH REPORT

Int. national Application No.

PCT/US 96/15524

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JOURNAL OF VIROLOGY, vol. 69, no. 9, September 1995, pages 5677-5686, XP000612069 GROSSELD H. ET AL.: "RNA replication by respiratory syncytial virus (RSV) is directed by the N, P, and L proteins; transcription also occurs under these conditions but requires RSV superinfection for efficient synthesis of full-length mRNA" cited in the application see page 5685, column 1, paragraph 6 - column 2, paragraph 2 ---	1,2
X	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 88, no. 21, November 1991, WASHINGTON US, pages 9663-9667, XP000196754 COLLINS P.L. ET AL.: "Rescue of synthetic analogs of respiratory syncytial virus genomic RNA and effect of truncations and mutations on the expression of a foreign reporter gene" cited in the application see the whole document ---	44-47
A	WO 93 21310 A (AMERICAN HOME PROD) 28 October 1993 ---	
A	WO 92 01471 A (THE UAB RESEARCH FOUNDATION) 6 February 1992 ---	
P,X	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 92, no. 25, 5 December 1995, WASHINGTON, US, pages 11563-11567, XP000196787 COLLINS P.L. ET AL.: "Production of infectious human respiratory syncytial virus from cloned cDNA confirms an essential role for the transcription elongation factor from the 5' proximal open reading frame of the M2 mRNA in gene expression and provides a capability for vaccine development" cited in the application see the whole document ---	1-47

-f--

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 96/15524

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>EMBO JOURNAL, vol. 13, no. 18, 1994, EYNSHAM, OXFORD GB, pages 4195-4203, XP000612065 SCHNELL M.J. ET AL.: "Infectious rabies viruses from cloned cDNA" cited in the application -----</p>	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/15524

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9207940	14-05-92	AU-A- 9128391	26-05-92
WO-A-9408022	14-04-94	AU-A- 5103893 EP-A- 0672153 JP-T- 8501453 ZA-A- 9307164	26-04-94 20-09-95 20-02-96 23-05-94
WO-A-9321310	28-10-93	AU-A- 4290793 CA-A- 2118509 EP-A- 0640128 HU-A- 69992 JP-T- 7505891 ZA-A- 9302766	18-11-93 28-10-93 01-03-95 28-09-95 29-06-95 20-10-94
WO-A-9201471	06-02-92	AU-B- 650040 AU-A- 8330391 CA-A- 2087853 EP-A- 0540645 HU-A- 67362 JP-T- 5509231 NZ-A- 239084 NZ-A- 250402	09-06-94 18-02-92 25-01-92 12-05-93 28-03-95 22-12-93 27-09-94 28-08-95